


*gluons  
measure  
size of  
color  
dipole*

$$\frac{d\sigma}{dk_t^2} \propto |\alpha_s(k_t^2) x_N G(u, k_t^2)|^2 \left| \frac{\partial^2}{\partial k_t^2} \psi(\mathbf{x}, k_t) \right|^2$$


# Key Ingredients in E791 Experiment



Brodsky Mueller  
Frankfurt Miller Strikman

*Small color-dipole moment pion not absorbed;  
interacts with each nucleon coherently*

## QCD COLOR Transparency



Diffraction 2008

Novel QCD Diffractive Physics

Stan Brodsky, SLAC


La Londe-les-Maures, September 13, 2008

- Fully coherent interactions between pion and nucleons.
- Emerging Di-Jets do not interact with nucleus.

$$M(A) = A \cdot M(N)$$

$$\frac{d\sigma}{dq_t^2} \propto A^2 \quad q_t^2 \sim 0$$

$$\sigma \propto A^{4/3}$$



# Measure pion LFWF in diffractive dijet production

## Confirmation of color transparency

A-Dependence results:  $\sigma \propto A^\alpha$

<u><math>k_t</math> range (GeV/c)</u>	<u><math>\alpha</math></u>	<u><math>\alpha</math> (CT)</u>
$1.25 < k_t < 1.5$	$1.64 +0.06 -0.12$	1.25
$1.5 < k_t < 2.0$	$1.52 \pm 0.12$	1.45
$2.0 < k_t < 2.5$	$1.55 \pm 0.16$	1.60

Ashery E791

$\alpha$  (Incoh.) =  $0.70 \pm 0.1$

*Conventional Glauber Theory Ruled  
Out!*

**Factor of 7**

Diffraction 2008

Novel QCD Diffractive Physics

Stan Brodsky, SLAC


La Londe-les-Maures, September 13, 2008

# Color Transparency

Bertsch, Gunion, Goldhaber, sjb  
A. H. Mueller, sjb

- Fundamental test of gauge theory in hadron physics
- Small color dipole moments interact weakly in nuclei
- Complete coherence at high energies
- Clear Demonstration of CT from Diffractive Di-Jets

# E791 Diffractive Di-Jet transverse momentum distribution



## Two Components

*High Transverse momentum dependence  $k_T^{-6.5}$  consistent with PQCD, ERBL Evolution*


*Gaussian component at small  $k_T$  similar to AdS/CFT LFWF*



# Diffractive Dissociation of a Pion into Dijets

$$\pi A \rightarrow \text{JetJet}A'$$

$$\Psi_{q\bar{q}}^{\pi}(x, \vec{k}_{\perp})$$

- E791 Fermilab Experiment  
Ashery et al
- 500 GeV pions collide on nuclei keeping it intact
- Measure momentum of two jets
- Study momentum distributions of pion LF wavefunction







Ashery E791

### *Narrowing of $x$ distribution at high jet transverse momentum*

$x$  distribution of diffractive dijets from the platinum target for  $1.25 \leq k_t \leq 1.5$  GeV/ $c$  (left) and for  $1.5 \leq k_t \leq 2.5$  GeV/ $c$  (right). The solid line is a fit to a combination of the asymptotic and CZ distribution amplitudes. The dashed line shows the contribution from the asymptotic function and the dotted line that of the CZ function.





Ashery  
E79I

**Possibly two components:**

**Perturbative (ERBL) + Nonperturbative (AdS/CFT)**

$$\phi(x) = A_{\text{pert}}(k_{\perp}^2)x(1-x) + B_{\text{nonpert}}(k_{\perp}^2)\sqrt{x(1-x)}$$

*Narrowing of  $x$  distribution at high jet transverse momentum*


**Diffraction 2008**

**Novel QCD Diffractive Physics**

**Stan Brodsky, SLAC**

**La Londe-les-Maures, September 13, 2008**

$$F_{\pi}(Q^2) = \int_0^1 dx \phi_{\pi}(x) \int_0^1 dy \phi_{\pi}(y) \frac{16\pi C_F \alpha_V(Q_V)}{(1-x)(1-y)Q^2}$$



$$\phi(x, Q_0) \propto \sqrt{x(1-x)}$$

$$\phi_{asymptotic} \propto x(1-x)$$

Normalized to  $f_{\pi}$

***AdS/CFT:***

Increases PQCD leading twist prediction for  $F_{\pi}(Q^2)$  by factor 16/9

Diffraction 2008


Novel QCD Diffractive Physics

La Londe-les-Maures, September 13, 2008

Stan Brodsky, SLAC

# Diffractive Dissociation of Proton into Quark Jets

Frankfurt, Miller,  
Strikman



Measure Light-Front Wavefunction of Proton

Minimal momentum transfer to nucleus  
Nucleus left Intact!

Diffraction 2008

Novel QCD Diffractive Physics

Stan Brodsky, SLAC


La Londe-les-Maures, September 13, 2008

# Electromagnetic Tri-Jet Excitation of Proton

$$ep \rightarrow e \text{ jet jet jet}$$

Measure light-front  
wavefunction of  
proton

$$\frac{\partial}{\partial k_{\perp}} \Psi_{n=3}^p(x_i, \vec{k}_{\perp i}, \lambda_i)$$




Need Forward  
Small Angle  
Detection


# Measurement of Charm Structure Function

J. J. Aubert et al. [European Muon Collaboration], "Production Of Charmed Particles In 250-GeV Mu+ - Iron Interactions," Nucl. Phys. B 213, 31 (1983).

## First Evidence for Intrinsic Charm




*factor of 30!*



***DGLAP / Photon-Gluon Fusion: factor of 30 too small***

- EMC data:  $c(x, Q^2) > 30 \times \text{DGLAP}$   
 $Q^2 = 75 \text{ GeV}^2, x = 0.42$
- High  $x_F$   $pp \rightarrow J/\psi X$
- High  $x_F$   $pp \rightarrow J/\psi J/\psi X$
- High  $x_F$   $pp \rightarrow \Lambda_c X$
- High  $x_F$   $pp \rightarrow \Lambda_b X$
- High  $x_F$   $pp \rightarrow \Xi(ccd) X$  (SELEX)

## IC Structure Function: Critical Test of QCD



$|uudc\bar{c}\rangle$  Fluctuation in Proton

QCD: Probability  $\sim \frac{\Lambda_{QCD}^2}{M_Q^2}$

$|e^+e^-l^+l^-\rangle$  Fluctuation in Positronium

QED: Probability  $\sim \frac{(m_e\alpha)^4}{M_l^4}$

OPE derivation - M.Polyakov et al.

$$\langle p | \frac{G_{\mu\nu}^3}{m_Q^2} | p \rangle \text{ vs. } \langle p | \frac{F_{\mu\nu}^4}{m_l^4} | p \rangle \quad c\bar{c} \text{ in Color Octet}$$

Distribution peaks at equal rapidity (velocity)  
Therefore heavy particles carry the largest momentum fractions


$$\hat{x}_i = \frac{m_{\perp i}}{\sum_j^n m_{\perp j}}$$

*High x charm!*

Hoyer, Peterson, Sakai, sjb

# Intrinsic Heavy-Quark Fock States


- Rigorous prediction of QCD, OPE
- Color-Octet Color-Octet Fock State!



- Probability  $P_{Q\bar{Q}} \propto \frac{1}{M_Q^2}$   $P_{Q\bar{Q}Q\bar{Q}} \sim \alpha_s^2 P_{Q\bar{Q}}$   $P_{c\bar{c}/p} \simeq 1\%$
- Large Effect at high x
- Greatly increases kinematics of colliders such as Higgs production (Kopeliovich, Schmidt, Soffer, sjb)
- Severely underestimated in conventional parameterizations of heavy quark distributions (Pumplin, Tung)
- Many empirical tests



# Leading Hadron Production from Intrinsic Charm





Coalescence of Comoving Charm and Valence Quarks  
Produce  $J/\psi$ ,  $\Lambda_c$  and other Charm Hadrons at High  $x_F$

# SELEX $\Lambda_c^+$ Studies – $p_T$ Dependence

- $\Lambda_c^+$  production by  $\Sigma^-$  vs  $x_F$  shows harder spectrum at low  $p_T$  - consistent with an intrinsic charm picture.

(Vogt, Brodsky and Hoyer, Nucl. Phys. B383,683 (1992))






# *Production of a Double-Charm Baryon*

**SELEX high  $x_F$**        $\langle x_F \rangle = 0.33$

# Production of Two Charmonia at High $x_F$



Diffraction 2008

La Londe-les-Maures, September 13, 2008

Novel QCD Diffractive Physics


80

Stan Brodsky, SLAC

All events have  $x_{\psi\psi}^F > 0.4$  !

**Excludes 'color drag' model**

$$\pi A \rightarrow J/\psi J/\psi X$$



Intrinsic charm contribution to double quarkonium hadroproduction <sup>\*</sup>

R. Vogt<sup>a</sup>, S.J. Brodsky<sup>b</sup>

The probability distribution for a general  $n$ -parton intrinsic  $c\bar{c}$  Fock state as a function of  $x$  and  $k_T$  written as

$$\frac{dP_{ic}}{\prod_{i=1}^n dx_i d^2k_{T,i}} = N_n \alpha_s^4 (M_{c\bar{c}}) \frac{\delta(\sum_{i=1}^n k_{T,i}) \delta(1 - \sum_{i=1}^n x_i)}{(m_h^2 - \sum_{i=1}^n (m_{T,i}^2/x_i))^2},$$

Fig. 3. The  $\psi\psi$  pair distributions are shown in (a) and (c) for the pion and proton projectiles. Similarly, the distributions of  $J/\psi$ 's from the pairs are shown in (b) and (d). Our calculations are compared with the  $\pi^- N$  data at 150 and 280 GeV/c [1]. The  $x_{\psi\psi}$  distributions are normalized to the number of pairs from both pion beams (a) and the number of pairs from the 400 GeV proton measurement (c). The number of single  $J/\psi$ 's is twice the number of pairs.

## NA3 Data

Diffraction 2008

Novel QCD Diffractive Physics

La Londe-les-Maures, September 13, 2008

81


Stan Brodsky, SLAC

# Excitation of Intrinsic Heavy Quarks in Proton

Amplitude maximal at small invariant mass, equal rapidity

$$x_i \sim \frac{m_{\perp i}}{\sum_j^n m_{\perp j}}$$

Produce forward, high  $x_F$   
 $\Upsilon(b\bar{b}), \Lambda_b(bud), B^+(\bar{b}u), B^0(\bar{b}d)$



Diffraction 2008

Novel QCD Diffractive Physics


Stan Brodsky, SLAC

La Londe-les-Maures, September 13, 2008

82

800 GeV p-A (FNAL)  $\sigma_A = \sigma_p * A^\alpha$   
 PRL 84, 3256 (2000); PRL 72, 2542 (1994)

$$\frac{d\sigma}{dx_F} (pA \rightarrow J/\psi X)$$





Remarkably Strong Nuclear Dependence for Fast Charmonium

Violation of PQCD Factorization!

Violation of factorization in charm hadroproduction.

P. Hoyer, M. Vanttinen (Helsinki U.), U. Sukhatme (Illinois U., Chicago) . HU-TFT-90-14, May 1990. 7pp.  
 Published in Phys.Lett.B246:217-220,1990

## PHENIX compared to lower energy measurements



*Huge  
"absorption"  
effect at  
large  $x_F$*



Klein, Vogt, PRL 91:142301, 2003  
Kopeliovich, NP A696:669, 2001

*Violates PQCD  
factorization!*

Hoyer, Sukhatme, Vanttinen

$$\frac{d\sigma}{dx_F} (pA \rightarrow J/\psi X)$$



J. Badier et al, NA3  
*Two Components*






$$\frac{d\sigma}{dx_F}(pA \rightarrow J/\psi X) = A^1 \frac{d\sigma_1}{dx_F} + A^{2/3} \frac{d\sigma_{2/3}}{dx_F}$$

$A^1$  component

*Identify with Fusion*

**Conventional PQCD  
subprocesses**

$$\frac{d\sigma_1}{dx_F}(\pi A \rightarrow J/\psi X)$$




**Diffraction 2008**


**NOVERQCD Diffractive Physics**

J. Badier et al, NA3

$$\frac{d\sigma}{dx_F}(pA \rightarrow J/\psi X) = A^1 \frac{d\sigma_1}{dx_F} + A^{2/3} \frac{d\sigma_{2/3}}{dx_F}$$



$A^{2/3}$  component



*Identify with IC  
High  $x_F$*

*Remarkably Flat  
Distribution*

**Excess beyond conventional PQCD subprocesses**

Diffraction 2008

Novel QCD Diffractive Physics

Stan Brodsky, SLAC


La Londe-les-Maures, September 13, 2008

*Color-Opaque IC Fock state  
interacts on nuclear front surface*

*Scattering on front-face nucleon produces color-singlet  $c\bar{c}$  pair*

*Octet-Octet IC Fock State*

*No absorption of  
small color-singlet*




$$\frac{d\sigma}{dx_F}(pA \rightarrow J/\psi X) = A^{2/3} \times \frac{d\sigma}{dx_F}(pN \rightarrow J/\psi X)$$

- IC Explains Anomalous  $\alpha(x_F)$  not  $\alpha(x_2)$  dependence of  $pA \rightarrow J/\psi X$   
(Mueller, Gunion, Tang, SJB)
- Color Octet IC Explains  $A^{2/3}$  behavior at high  $x_F$  (NA3, Fermilab) *Color Opacity*  
(Kopeliovitch, Schmidt, Soffer, SJB)
- IC Explains  $J/\psi \rightarrow \rho\pi$  puzzle  
(Karliner, SJB)
- IC leads to new effects in  $B$  decay  
(Gardner, SJB)

## Higgs production at $x_F = 0.8$

# Measure $c(x)$ in Deep Inelastic Lepton-Proton Scattering



Hoyer, Peterson, SJB

## *Why is Intrinsic Charm Important for Flavor Physics?*

- New perspective on fundamental nonperturbative hadron structure
- Charm structure function at high  $x$
- Dominates high  $x_F$  charm and charmonium production
- Hadroproduction of new heavy quark states such as  $ccu$ ,  $ccd$  at high  $x_F$
- Intrinsic charm -- long distance contribution to penguin mechanisms for weak decay
- Novel Nuclear Effects from color structure of IC, Heavy Ion Collisions
- New mechanisms for high  $x_F$  Higgs hadroproduction
- Dynamics of  $b$  production: LHCb
- Fixed target program at LHC: produce  $bbb$  states


PHYSICAL REVIEW D **73**, 113005 (2006)

## **Diffraction Higgs production from intrinsic heavy flavors in the proton**

Stanley J. Brodsky,<sup>1,\*</sup> Boris Kopeliovich,<sup>2,†</sup> Ivan Schmidt,<sup>2,‡</sup> and Jacques Soffer<sup>3,§</sup>


<sup>1</sup>

$$\frac{d\sigma(pp \rightarrow ppH)}{dx_2 d^2 p_1 d^2 p_2} = \frac{1}{(1-x_2)16\pi^2} |A(x_2, \vec{p}_1, \vec{p}_2)|^2$$





$$\begin{aligned}
A(x_2, \vec{p}_1, \vec{p}_2) &= \frac{8}{3\sqrt{2}} \int d^2 Q \frac{d^2 q}{q^2} \frac{d^2 k}{k^2} \alpha_s(q^2) \alpha_s(k^2) \delta(\vec{q} + \vec{p}_2 + \vec{k}) \delta(\vec{k} - \vec{p}_1 - \vec{Q}) \\
&\times \int d^2 \tau |\Phi_p(\tau)|^2 [e^{i(\vec{k} + \vec{q}) \cdot \vec{\tau}/2} - e^{i(\vec{q} - \vec{k}) \cdot \vec{\tau}/2}] \\
&\times \int d^2 R d^2 r d^2 \rho H^\dagger(\vec{r}) e^{i\vec{q} \cdot \vec{r}/2} (1 - e^{-i\vec{q} \cdot \vec{r}}) \Phi_p^\dagger(\vec{\rho}) e^{i\vec{k} \cdot \vec{\rho}/2} (1 - e^{-i\vec{k} \cdot \vec{\rho}}) \Psi_p(\vec{R}, \vec{r}, \vec{\rho}, z) e^{i\vec{Q} \cdot \vec{R}}.
\end{aligned}$$




$$\Psi_p(\vec{R}, \vec{r}, \vec{\rho}, z) = \Psi_{\text{IC}}(\vec{R}, z) \Psi_{\bar{c}c}(\vec{r}) \Psi_{3q}(\vec{\rho}).$$


$$\int_0^1 dz \int d^2R d^2r d^2\rho |\Psi_p(\vec{R}, \vec{r}, \vec{\rho}, z)|^2 = P_{\text{IC}},$$


$$\Psi_{\text{IQ}}(Q, z, \kappa) \propto \frac{z(1-z)}{Q^2 + z^2 m_p^2 + M_{\bar{Q}Q}^2 (1-z)}.$$

$$H(\vec{r}) = i \frac{\sqrt{N_c G_F}}{2\pi} m_c \bar{\chi} \vec{\sigma} \chi \frac{\vec{r}}{r} \left[ \epsilon Y_1(\epsilon r) - \frac{ir}{2} \Gamma_H M_H Y_0(\epsilon r) \right]$$




The distribution of produced Higgs particles over the fraction of the proton beam momentum. The dotted, dashed, and solid curves correspond to Higgs production from nonperturbative IC ( $\beta = 1$ ), perturbative IC ( $\beta = 0$ ), and IT, respectively.






The cross section of the reaction  $pp \rightarrow Hp + p$  as a function of the Higgs mass. Contributions of IC (dashed line), IB (dotted line), and IT (solid line).

$J/\Psi, \chi_c, Y, \chi_b, Z^0, H$



*Particle ratio changes with centrality!*



*Protons less absorbed  
in nuclear collisions than pions*


← **Central**

- ■ Au+Au 0-10%
- △ ▲ Au+Au 20-30%
- ● Au+Au 60-92%
- ★ p+p,  $\sqrt{s} = 53$  GeV, ISR
- e<sup>+</sup>e<sup>-</sup>, gluon jets, DELPHI
- ..... e<sup>+</sup>e<sup>-</sup>, quark jets, DELPHI

← **Peripheral**

$pp \rightarrow HX$  at high  $p_T$   
*Proton created from  
jet fragmentation*

*Color Opaque*



$$n_{\text{active}} = 4$$

$$n = 2n_{\text{active}} - 4$$

$$n = 4$$

**u**  
**Novel QCD Diffractive Physics**

**Diffractive 2008**

**La Londe-les-Maures, September 13, 2008**

**Stan Brodsky, SLAC**

# Crucial Test of Leading -Twist QCD: Scaling at fixed $x_T$

$$x_T = \frac{2p_T}{\sqrt{s}}$$

$$E \frac{d\sigma}{d^3p}(pN \rightarrow \pi X) = \frac{F(x_T, \theta_{CM})}{p_T^{n_{eff}}}$$


**Parton model:  $n_{eff} = 4$**

***As fundamental as Bjorken scaling in DIS***

**Conformal scaling:  $n_{eff} = 2 n_{active} - 4$**



QCD prediction: Modification of power fall-off due to DGLAP evolution and the Running Coupling




*Key test of PQCD: power-law fall-off at fixed x<sub>T</sub>*


$$pp \rightarrow \gamma X$$

$$E \frac{d\sigma}{d^3p}(pp \rightarrow \gamma X) = \frac{F(\theta_{cm}, x_T)}{p_T^4}$$

$$gu \rightarrow \gamma u$$




$$\sqrt{s}^n E \frac{d\sigma}{d^3p} (pp \rightarrow \gamma X) \text{ at fixed } x_T$$



**Scaling of direct  
photon  
production  
consistent with  
PQCD**

$$\sqrt{s}^{6.3} \times E \frac{d\sigma}{d^3p} (pp \rightarrow H^\pm X) \text{ at fixed } x_T$$



Tannenbaum

**Scaling  
inconsistent with  
PQCD**


Diffraction 2008

Novel QCD Diffractive Physics

La Londe-les-Maures, September 13, 2008


Stan Brodsky, SLAC

$$E \frac{d\sigma}{d^3p} (pp \rightarrow HX) = \frac{F(x_T, \theta_{cm} = \pi/2)}{p_T^n}$$



*Clear evidence for higher-twist contributions*

**J. W. Cronin, SSI 1974**



**Diffraction 2008**


**Novel QCD Diffractive Physics**

**La Londe-les-Maures, September 13, 2008**

**105**

**Stan Brodsky, SLAC**

Protons produced in AuAu collisions at RHIC do not exhibit clear scaling properties in the available  $p_T$  range. Shown are data for central (0 – 5%) and for peripheral (60 – 90%) collisions.



Diffraction 2008



Novel QCD Diffractive Physics

Stan Brodsky, SLAC

La Londe-les-Maures, September 13, 2008

106

$$E \frac{d\sigma}{d^3p} (pp \rightarrow HX) = \frac{F(x_T, \theta_{CM})}{n_{eff} p_T}$$



$$E \frac{d\sigma}{d^3p} (pp \rightarrow pX) = \frac{F(x_T, \theta_{CM})}{p_T^{12}}$$

$$E \frac{d\sigma}{d^3p} (pp \rightarrow pX) = \frac{F(x_T, \theta_{CM})}{p_T^8}$$

*Trend consistent with RHIC at small  $x_T$*

Diffraction 2008

$x_T$

Small QCD Diffractive Physics

La Londe-les-Maures, September 13, 2008

107

Stan Brodsky, SLAC

# Baryon can be made directly within hard subprocess

## Coalescence within hard subprocess

$$b_{\perp} \simeq 1/p_T$$

Bjorken

Blankenbecler, Gunion, sjb


Berger, sjb

Hoyer, et al: Semi-Exclusive

$$uu \rightarrow p\bar{d}$$

$$\phi_p(x_1, x_2, x_3) \propto \Lambda_{QCD}^2$$

*Small color-singlet  
Color Transparent  
Minimal same-side energy*



*Collision can produce 3 collinear quarks*

$$n_{\text{active}} = 6$$

$$n_{\text{eff}} = 2n_{\text{active}} - 4$$

$$n_{\text{eff}} = 8$$

Diffraction 2008

Novel QCD Diffractive Physics


Stan Brodsky, SLAC

La Londe-les-Maures, September 13, 2008


108



*Baryon made directly within hard subprocess*



$$\sqrt{s_{NN}} = 130 \text{ and } 200 \text{ GeV}$$



*Proton power changes with centrality !*

Diffraction 2008

Novel QCD Diffractive Physics


Stan Brodsky, SLAC

La Londe-les-Maures, September 13, 2008

IIO

Power-law exponent  $n(x_T)$  for  $\pi^0$  and  $h$  spectra in central and peripheral Au+Au collisions at  $\sqrt{s_{NN}} = 130$  and 200 GeV

S. S. Adler, *et al.*, PHENIX Collaboration, *Phys. Rev. C* **69**, 034910 (2004) [nucl-ex/0308006].



*Proton power changes with centrality !*

*Proton production dominated by color-transparent direct high  $n_{eff}$  subprocesses*

Diffraction 2008


Novel QCD Diffractive Physics

Stan Brodsky, SLAC

La Londe-les-Maures, September 13, 2008

III

*Particle ratio changes with centrality!*



*Protons less absorbed  
in nuclear collisions than pions  
because of dominant  
color transparent higher twist process*

← **Central**

- ■ Au+Au 0-10%
- △ ▲ Au+Au 20-30%
- ● Au+Au 60-92%
- ★ p+p,  $\sqrt{s} = 53$  GeV, ISR
- e<sup>+</sup>e<sup>-</sup>, gluon jets, DELPHI
- ..... e<sup>+</sup>e<sup>-</sup>, quark jets, DELPHI

← **Peripheral**


Diffraction 2008

Novel QCD Diffractive Physics

La Londe-les-Maures, September 13, 2008

II2

Stan Brodsky, SLAC




*proton trigger:  
# same-side particles  
decreases with centrality*



**Proton production more dominated by color-transparent direct high- $n_{eff}$  subprocesses**

Baryon to Meson Ratios



Diffraction 2008

Novel QCD Diffractive Physics

Stan Brodsky, SLAC

La Londe-les-Maures, September 13, 2008


114

*Lambda can be made directly within hard subprocess*

**Coalescence  
within hard  
subprocess**

$$ud \rightarrow \Lambda \bar{s}$$

*Small color-singlet  
Color Transparent  
Minimal same-side energy*




# *Baryon Anomaly: Evidence for Direct, Higher-Twist Subprocesses*

- **Explains anomalous power behavior at fixed  $x_T$**
- **Protons more likely to come from direct higher-twist subprocess than pions**
- **Protons less absorbed than pions in central nuclear collisions because of color transparency**
- **Predicts increasing proton to pion ratio in central collisions**
- **Proton power  $n_{\text{eff}}$  increases with centrality since leading twist contribution absorbed**
- **Fewer same-side hadrons for proton trigger at high centrality**
- **Exclusive-inclusive connection at  $x_T = 1$**




- Quarks and Gluons:  
Fundamental constituents of hadrons and nuclei



- *Quantum Chromodynamics (QCD)*

- New Insights from higher space-time dimensions: *AdS/QCD*


- *Light-Front Holography*



- *Hadronization at the Amplitude Level*

- *Light Front Wavefunctions:* analogous to the Schrodinger wavefunctions of atomic physics

$$\Psi_n(x_i, \vec{k}_{\perp i}, \lambda_i)$$



# Novel Aspects of QCD

- Heavy quark distributions **do not** derive exclusively from DGLAP or gluon splitting -- **component intrinsic to hadron wavefunction: Higgs at high  $x_F$**
- Initial and final-state interactions **are not** power suppressed in hard QCD reactions
- LFWFS are universal, but measured nuclear parton distributions **are not** universal -- **antishadowing is flavor dependent**
- Hadroproduction at large transverse momentum **does not** derive exclusively from 2 to 2 scattering subprocesses

- **DDIS and Sivers Effect: Breakdown of Leading-Twist Factorization**
- **Physics of Hard Pomeron**
- **Measure Fundamental Hadron Wavefunction via Di-jet and Tri-jet Fragmentation**
- **Origin of Leading Twist Shadowing**
- **Non-Universal Antishadowing**
- **Heavy quark structure functions at high  $x$**
- **Higgs production at large  $x_F$**
- **Hadroproduction of new heavy quark states such as  $ccu$ ,  $ccd$  at high  $x_F$**
- **Novel Nuclear Effects from color structure of IC**
- **Fixed target program at LHC: produce  $bbb$  states**
- **Direct Hadroproduction at high  $p_T$**