


Light-Front Holography and Gauge/Gravity Correspondence: Applications to the Meson and Baryon Spectrum

Guy F. de Téramond

University of Costa Rica

In Collaboration with Stan Brodsky



Light-Cone 2009: Relativistic Hadronic and Particle Physics

Instituto Tecnológico de Aeronáutica

São José dos Campos, Brazil, July 8-13, 2009



GdT and Brodsky, PRL **102**, 081601 (2009)

Outline


1. Introduction
2. Light-Front Dynamics
 - Light-Front Fock Representation
3. Semiclassical Approximation to QCD
 - Hard-Wall Model
 - Holographic Mapping
4. Higher-Spin Bosonic Modes
 - Hard-Wall Model
 - Soft-Wall Model
5. Higher-Spin Fermionic Modes
 - Hard-Wall Model
 - Soft-Wall Model

1 Introduction

- Most challenging problem of strong interaction dynamics: determine the composition of hadrons in terms of their fundamental QCD quark and gluon degrees of freedom
- Recent developments inspired by the AdS/CFT correspondence [Maldacena (1998)] between string states in AdS space and conformal field theories in physical space-time have led to analytical insights into the confining dynamics of QCD
- Description of strongly coupled gauge theory using a dual gravity description!
- Strings describe spin- J extended objects (no quarks). QCD degrees of freedom are pointlike particles and hadrons have orbital angular momentum: how can they be related?
- Light-front (LF) quantization is the ideal framework to describe hadronic structure in terms of quarks and gluons: simple vacuum structure allows unambiguous definition of the partonic content of a hadron, exact formulae for form factors, physics of angular momentum of constituents ...
- Frame-independent LF Hamiltonian equation $P_\mu P^\mu |P\rangle = \mathcal{M}^2 |P\rangle$ similar structure of AdS EOM
- First semiclassical approximation to the bound-state LF Hamiltonian equation in QCD is equivalent to equations of motion in AdS and can be systematically improved

2 Light Front Dynamics

- Different possibilities to parametrize space-time [Dirac (1949)]
- Parametrizations differ by the hypersurface on which the initial conditions are specified. Each evolve with different “times” and has its own Hamiltonian, but should give the same physical results
- *Instant form*: hypersurface defined by $t = 0$, the familiar one
- *Front form*: hypersurface is tangent to the light cone at $\tau = t + z/c = 0$



$$x^+ = x^0 + x^3 \quad \text{light-front time}$$


$$x^- = x^0 - x^3 \quad \text{longitudinal space variable}$$

$$k^+ = k^0 + k^3 \quad \text{longitudinal momentum} \quad (k^+ > 0)$$

$$k^- = k^0 - k^3 \quad \text{light-front energy}$$

$$k \cdot x = \frac{1}{2} (k^+ x^- + k^- x^+) - \mathbf{k}_\perp \cdot \mathbf{x}_\perp$$

On shell relation $k^2 = m^2$ leads to dispersion relation $k^- = \frac{\mathbf{k}_\perp^2 + m^2}{k^+}$



- QCD Lagrangian

$$\mathcal{L}_{\text{QCD}} = -\frac{1}{4g^2} \text{Tr} (G^{\mu\nu} G_{\mu\nu}) + i\bar{\psi} D_\mu \gamma^\mu \psi + m\bar{\psi}\psi$$

- LF Momentum Generators $P = (P^+, P^-, \mathbf{P}_\perp)$ in terms of dynamical fields ψ, \mathbf{A}_\perp

$$P^- = \frac{1}{2} \int dx^- d^2 \mathbf{x}_\perp \bar{\psi} \gamma^+ \frac{(i\nabla_\perp)^2 + m^2}{i\partial^+} \psi + \text{interactions}$$

$$P^+ = \int dx^- d^2 \mathbf{x}_\perp \bar{\psi} \gamma^+ i\partial^+ \psi$$


$$\mathbf{P}_\perp = \frac{1}{2} \int dx^- d^2 \mathbf{x}_\perp \bar{\psi} \gamma^+ i\nabla_\perp \psi$$

- LF Hamiltonian P^- generates LF time translations

$$[\psi(x), P^-] = i \frac{\partial}{\partial x^+} \psi(x)$$

and the generators P^+ and \mathbf{P}_\perp are kinematical

Light-Front Fock Representation



- Dirac field ψ , expanded in terms of ladder operators on the initial surface $x^+ = x^0 + x^3$

$$P^- = \sum_{\lambda} \int \frac{dq^+ d^2 \mathbf{q}_{\perp}}{(2\pi)^3} \left(\frac{\mathbf{q}_{\perp}^2 + m^2}{q^+} \right) b_{\lambda}^{\dagger}(q) b_{\lambda}(q) + \text{interactions}$$

Sum over free quanta $q^- = \frac{\mathbf{q}_{\perp}^2 + m^2}{q^+}$ plus interactions ($m^2 = 0$ for gluons)

- Construct light-front invariant Hamiltonian for the composite system: $H_{LF} = P_{\mu} P^{\mu} = P^- P^+ - \mathbf{P}_{\perp}^2$

$$H_{LC} |\psi_H\rangle = \mathcal{M}_H^2 |\psi_H\rangle$$

- State $|\psi_H(P)\rangle = |\psi_H(P^+, \mathbf{P}_{\perp}, J_z)\rangle$ is an expansion in multi-particle Fock eigenstates $|n\rangle$ of the free LF Hamiltonian:

$$|\psi_H\rangle = \sum_n \psi_{n/H} |n\rangle$$

- Fock components $\psi_{n/H}(x_i, \mathbf{k}_{\perp i}, \lambda_i^z)$ are independent of P^+ and \mathbf{P}_{\perp} and depend only on relative partonic coordinates: momentum fraction $x_i = k_i^+ / P^+$, transverse momentum $\mathbf{k}_{\perp i}$ and spin λ_i^z

$$\sum_{i=1}^n x_i = 1, \quad \sum_{i=1}^n \mathbf{k}_{\perp i} = 0.$$

- Compute \mathcal{M}^2 from hadronic matrix element

$$\langle \psi_H(P') | H_{LF} | \psi_H(P) \rangle = \mathcal{M}_H^2 \langle \psi_H(P') | \psi_H(P) \rangle$$

- Find

$$\mathcal{M}_H^2 = \sum_n \int [dx_i] [d^2\mathbf{k}_{\perp i}] \sum_{\ell} \left(\frac{\mathbf{k}_{\perp \ell}^2 + m_{\ell}^2}{x_{\ell}} \right) |\psi_{n/H}(x_i, \mathbf{k}_{\perp i})|^2 + \text{interactions}$$

- Phase space normalization of LFWFs

$$\sum_n \int [dx_i] [d^2\mathbf{k}_{\perp i}] |\psi_{n/h}(x_i, \mathbf{k}_{\perp i})|^2 = 1$$


- In terms of $n-1$ independent transverse impact coordinates $\mathbf{b}_{\perp j}, j = 1, 2, \dots, n-1,$

$$\mathcal{M}_H^2 = \sum_n \prod_{j=1}^{n-1} \int dx_j d^2\mathbf{b}_{\perp j} \psi_{n/H}^*(x_i, \mathbf{b}_{\perp i}) \sum_{\ell} \left(\frac{-\nabla_{\mathbf{b}_{\perp \ell}}^2 + m_{\ell}^2}{x_{\ell}} \right) \psi_{n/H}(x_i, \mathbf{b}_{\perp i}) + \text{interactions}$$

- Normalization

$$\sum_n \prod_{j=1}^{n-1} \int dx_j d^2\mathbf{b}_{\perp j} |\psi_n(x_j, \mathbf{b}_{\perp j})|^2 = 1$$

3 Semiclassical Approximation to QCD



- Consider a two-parton hadronic bound state in transverse impact space in the limit $m_q \rightarrow 0$

$$\mathcal{M}^2 = \int_0^1 \frac{dx}{1-x} \int d^2 \mathbf{b}_\perp \psi^*(x, \mathbf{b}_\perp) (-\nabla_{\mathbf{b}_\perp}^2) \psi(x, \mathbf{b}_\perp) + \text{interactions}$$

- Separate angular, transverse and longitudinal modes in terms of boost invariant transverse variable:

$$\zeta^2 = x(1-x) \mathbf{b}_\perp^2 \quad - \text{ In } \mathbf{k}_\perp \text{ space key variable is the LF KE } \mathbf{k}_\perp^2 / x(1-x) \quad -$$

$$\psi(x, \zeta, \varphi) = \frac{\phi(\zeta)}{\sqrt{2\pi\zeta}} e^{iM\varphi} X(x)$$

- Find ($L = |M|$)

$$\mathcal{M}^2 = \int d\zeta \phi^*(\zeta) \sqrt{\zeta} \left(-\frac{d^2}{d\zeta^2} - \frac{1}{\zeta} \frac{d}{d\zeta} + \frac{L^2}{\zeta^2} \right) \frac{\phi(\zeta)}{\sqrt{\zeta}} + \int d\zeta \phi^*(\zeta) U(\zeta) \phi(\zeta)$$

where the confining forces from the interaction terms is summed up in the effective potential $U(\zeta)$

- Ultra relativistic limit $m_q \rightarrow 0$ longitudinal modes $X(x)$ decouple and LF eigenvalue equation $H_{LF}|\phi\rangle = \mathcal{M}^2|\phi\rangle$ is a LF wave equation for ϕ

$$\left(\underbrace{-\frac{d^2}{d\zeta^2} - \frac{1-4L^2}{4\zeta^2}}_{\text{kinetic energy of partons}} + \underbrace{U(\zeta)}_{\text{confinement}} \right) \phi(\zeta) = \mathcal{M}^2 \phi(\zeta)$$



- Effective light-front Schrödinger equation: relativistic, frame-independent and analytically tractable
- Eigenmodes $\phi(\zeta)$ determine the hadronic mass spectrum and represent the probability amplitude to find n -massless partons at transverse impact separation ζ within the hadron at equal light-front time
- LF modes $\phi(\zeta) = \langle \zeta | \phi \rangle$ are normalized by

$$\langle \phi | \phi \rangle = \int d\zeta |\langle \zeta | \phi \rangle|^2 = 1$$

- Semiclassical approximation to light-front QCD does not account for particle creation and absorption but can be implemented in the LF Hamiltonian EOM

Hard-Wall Model

- Consider the potential (hard wall)

$$U(\zeta) = \begin{cases} 0 & \text{if } \zeta \leq \frac{1}{\Lambda_{\text{QCD}}} \\ \infty & \text{if } \zeta > \frac{1}{\Lambda_{\text{QCD}}} \end{cases}$$

- If $L^2 \geq 0$ the Hamiltonian is positive definite $\langle \phi | H_{LF}^L | \phi \rangle \geq 0$ and thus $\mathcal{M}^2 \geq 0$
- If $L^2 < 0$ the Hamiltonian is not bounded from below (“Fall-to-the-center” problem in Q.M.)
- Critical value of the potential corresponds to $L = 0$, the lowest possible stable state
- Solutions:

$$\phi_L(\zeta) = C_L \sqrt{\zeta} J_L(\zeta \mathcal{M})$$


- Mode spectrum from boundary conditions

$$\phi\left(\zeta = \frac{1}{\Lambda_{\text{QCD}}}\right) = 0$$

Thus


$$\mathcal{M}^2 = \beta_{Lk} \Lambda_{\text{QCD}}$$

- Excitation spectrum hard-wall model: $\mathcal{M}_{n,L} \sim L + 2n$



Light-meson orbital spectrum $\Lambda_{QCD} = 0.32$ GeV

Holographic Mapping



- Holographic mapping found originally by matching expressions of EM and gravitational form factors of hadrons in AdS and LF QCD [Brodsky and GdT (2006, 2008)]

- Substitute $\Phi(\zeta) \sim \zeta^{3/2}\phi(\zeta)$, $\zeta \rightarrow z$ in the conformal LFWE

$$\left(-\frac{d^2}{d\zeta^2} - \frac{1 - 4L^2}{4\zeta^2} \right) \phi(\zeta) = \mathcal{M}^2 \phi(\zeta)$$

- Find:

$$[z^2 \partial_z^2 - 3z \partial_z + z^2 \mathcal{M}^2 - (\mu R)^2] \Phi(z) = 0$$

with $(\mu R)^2 = -4 + L^2$, the wave equation of string mode in AdS₅ !

- Isomorphism of $SO(4, 2)$ group of conformal QCD with generators $P^\mu, M^{\mu\nu}, D, K^\mu$ with the group of isometries of AdS₅ space: $x^\mu \rightarrow \lambda x^\mu, z \rightarrow \lambda z$

$$ds^2 = \frac{R^2}{z^2} (\eta_{\mu\nu} dx^\mu dx^\nu - dz^2)$$


- AdS Breitenlohner-Freedman bound $(\mu R)^2 \geq -4$ equivalent to LF QM stability condition $L^2 \geq 0$
- Conformal dimension Δ of AdS mode Φ given in terms of 5-dim mass by $(\mu R)^2 = \Delta(\Delta - 4)$. Thus $\Delta = 2 + L$ in agreement with the twist scaling dimension of a two parton object in QCD

- AdS₅ metric:

$$\underbrace{ds^2}_{L_{\text{AdS}}} = \frac{R^2}{z^2} \underbrace{(\eta_{\mu\nu} dx^\mu dx^\nu - dz^2)}_{L_{\text{Minkowski}}}$$

- A distance L_{AdS} shrinks by a warp factor as observed in Minkowski space ($dz = 0$):

$$L_{\text{Minkowski}} \sim \frac{z}{R} L_{\text{AdS}}$$



- Different values of z correspond to different scales at which the hadron is examined
- Since $x^\mu \rightarrow \lambda x^\mu$, $z \rightarrow \lambda z$, short distances $x_\mu x^\mu \rightarrow 0$ maps to UV conformal AdS₅ boundary $z \rightarrow 0$, which corresponds to the $Q \rightarrow \infty$ UV zero separation limit
- Large confinement dimensions $x_\mu x^\mu \sim 1/\Lambda_{\text{QCD}}^2$ maps to large IR region of AdS₅, $z \sim 1/\Lambda_{\text{QCD}}$, thus there is a maximum separation of quarks and a maximum value of z at the IR boundary
- Local operators like \mathcal{O} and \mathcal{L}_{QCD} defined in terms of quark and gluon fields at the AdS₅ boundary
- Use the isometries of AdS to map the local interpolating operators at the UV boundary of AdS into the modes propagating inside AdS

4 Higher-Spin Bosonic Modes

Hard-Wall Model

- AdS_{d+1} metric $x^\ell = (x^\mu, z)$:

$$ds^2 = g_{\ell m} dx^\ell dx^m = \frac{R^2}{z^2} (\eta_{\mu\nu} dx^\mu dx^\nu - dz^2)$$

- Action for gravity coupled to scalar field in AdS_{d+1}

$$S = \int d^{d+1}x \sqrt{g} \left(\underbrace{\frac{1}{\kappa^2} (\mathcal{R} - 2\Lambda)}_{S_G} + \underbrace{\frac{1}{2} (g^{\ell m} \partial_\ell \Phi \partial_m \Phi - \mu^2 \Phi^2)}_{S_M} \right)$$

- Equations of motion for S_M

$$z^3 \partial_z \left(\frac{1}{z^3} \partial_z \Phi \right) - \partial_\rho \partial^\rho \Phi - \left(\frac{\mu R}{z} \right)^2 \Phi = 0$$

- Physical AdS modes $\Phi_P(x, z) \sim e^{-iP \cdot x} \Phi(z)$ are plane waves along the Poincaré coordinates with four-momentum P^μ and hadronic invariant mass states $P_\mu P^\mu = \mathcal{M}^2$
- Factoring out dependence of string mode $\Phi_P(x, z)$ along x^μ -coordinates

$$\left[z^2 \partial_z^2 - (d-1)z \partial_z + z^2 \mathcal{M}^2 - (\mu R)^2 \right] \Phi(z) = 0$$

- Spin J -field on AdS represented by rank- J totally symmetric tensor field $\Phi(x, z)_{\ell_1 \dots \ell_J}$ [Fronsdal; Fradkin and Vasiliev]

- Action in AdS_{d+1} for spin- J field

$$S_M = \frac{1}{2} \int d^{d+1}x \sqrt{g} \left(\partial_\ell \Phi_{\ell_1 \dots \ell_J} \partial^\ell \Phi^{\ell_1 \dots \ell_J} - \mu^2 \Phi_{\ell_1 \dots \ell_J} \Phi^{\ell_1 \dots \ell_J} + \dots \right)$$

- Each hadronic state of total spin J is dual to a normalizable string mode

$$\Phi_P(x, z)_{\mu_1 \dots \mu_J} = e^{-iP \cdot x} \Phi(z)_{\mu_1 \dots \mu_J}$$

with four-momentum P_μ , spin polarization indices along the 3+1 physical coordinates and hadronic invariant mass $P_\mu P^\mu = \mathcal{M}^2$

- For string modes with all indices along Poincaré coordinates, $\Phi_{z\mu_2 \dots \mu_J} = \Phi_{\mu_1 z \dots \mu_J} = \dots = 0$ and appropriate subsidiary conditions system of coupled differential equations from S_M reduce to a homogeneous wave equation for $\Phi(z)_{\mu_1 \dots \mu_J}$

- Obtain spin- J mode $\Phi_{\mu_1 \dots \mu_J}$ with all indices along 3+1 coordinates from Φ by shifting dimensions

$$\Phi_J(z) = \left(\frac{z}{R}\right)^{-J} \Phi(z)$$

- Normalization [Hong, Yoon and Strassler (2006)]

$$R^{d-2J-1} \int_0^{z_{max}} \frac{dz}{z^{d-2J-1}} \Phi_J^2(z) = 1$$

- Substituting in the AdS scalar wave equation for Φ

$$\left[z^2 \partial_z^2 - (d-1-2J)z \partial_z + z^2 \mathcal{M}^2 - (\mu R)^2 \right] \Phi_J = 0$$

upon fifth-dimensional mass rescaling $(\mu R)^2 \rightarrow (\mu R)^2 - J(d-J)$

- Conformal dimension of J -mode

$$\Delta = \frac{1}{2} \left(d + \sqrt{(d-2J)^2 + 4\mu^2 R^2} \right)$$

and thus $(\mu R)^2 = (\Delta - J)(\Delta - d + J)$

- Upon substitution $z \rightarrow \zeta$ and

$$\phi_J(\zeta) \sim \zeta^{-3/2+J} \Phi_J(\zeta)$$

we recover the QCD LF wave equation ($d = 4$)

$$\left(-\frac{d^2}{d\zeta^2} - \frac{1 - 4L^2}{4\zeta^2} \right) \phi_{\mu_1 \dots \mu_J} = \mathcal{M}^2 \phi_{\mu_1 \dots \mu_J}$$



with $(\mu R)^2 = -(2 - J)^2 + L^2$

- J -decoupling in the HW model
- For $L^2 \geq 0$ the LF Hamiltonian is positive definite $\langle \phi_J | H_{LF} | \phi_J \rangle \geq 0$ and we find the stability bound $(\mu R)^2 \geq -(2 - J)^2$
- The scaling dimensions are $\Delta = 2 + L$ independent of J in agreement with the twist scaling dimension of a two parton bound state in QCD

Soft-Wall Model



- Soft-wall model [Karch, Katz, Son and Stephanov (2006)] retain conformal AdS metrics but introduce smooth cutoff which depends on the profile of a dilaton background field $\varphi(z) = \pm \kappa^2 z^2$

$$S = \int d^d x dz \sqrt{g} e^{\varphi(z)} \mathcal{L},$$

- Equation of motion for scalar field $\mathcal{L} = \frac{1}{2} (g^{\ell m} \partial_\ell \Phi \partial_m \Phi - \mu^2 \Phi^2)$

$$[z^2 \partial_z^2 - (d - 1 \mp 2\kappa^2 z^2) z \partial_z + z^2 \mathcal{M}^2 - (\mu R)^2] \Phi(z) = 0$$

with $(\mu R)^2 \geq -4$.

- LH holography requires 'plus dilaton' $\varphi = +\kappa^2 z^2$. Lowest possible state $(\mu R)^2 = -4$

$$\mathcal{M}^2 = 0, \quad \Phi(z) \sim z^2 e^{-\kappa^2 z^2}, \quad \langle r^2 \rangle \sim \frac{1}{\kappa^2}$$

A chiral symmetric bound state of two massless quarks with scaling dimension 2: the pion

- Action in AdS_{d+1} for spin J -field

$$S_M = \frac{1}{2} \int d^d x dz \sqrt{g} e^{\kappa^2 z^2} \left(\partial_{\ell} \Phi_{\ell_1 \dots \ell_J} \partial^{\ell} \Phi^{\ell_1 \dots \ell_J} - \mu^2 \Phi_{\ell_1 \dots \ell_J} \Phi^{\ell_1 \dots \ell_J} + \dots \right)$$

- Obtain spin- J mode $\Phi_{\mu_1 \dots \mu_J}$ with all indices along 3+1 coordinates from Φ by shifting dimensions

$$\Phi_J(z) = \left(\frac{z}{R} \right)^{-J} \Phi(z)$$

- Normalization

$$R^{d-2J-1} \int_0^{\infty} \frac{dz}{z^{d-2J-1}} e^{\kappa^2 z^2} \Phi_J^2(z) = 1.$$

- Substituting in the AdS scalar wave equation for Φ

$$\left[z^2 \partial_z^2 - (d-1-2J-2\kappa^2 z^2) z \partial_z + z^2 \mathcal{M}^2 - (\mu R)^2 \right] \Phi_J = 0$$

upon mass rescaling $(\mu R)^2 \rightarrow (\mu R)^2 - J(d-J)$ and $\mathcal{M}^2 \rightarrow \mathcal{M}^2 - 2J\kappa^2$

- Upon substitution $z \rightarrow \zeta$ ($J_z = L_z + S_z$) we find for $d = 4$

$$\phi_J(\zeta) \sim \zeta^{-3/2+J} e^{\kappa^2 \zeta^2 / 2} \Phi_J(\zeta), \quad (\mu R)^2 = -(2 - J)^2 + L^2$$

$$\left(-\frac{d^2}{d\zeta^2} - \frac{1 - 4L^2}{4\zeta^2} + \kappa^4 \zeta^2 + 2\kappa^2(L + S - 1) \right) \phi_{\mu_1 \dots \mu_J} = \mathcal{M}^2 \phi_{\mu_1 \dots \mu_J}$$



- Eigenfunctions

$$\phi_{nL}(\zeta) = \kappa^{1+L} \sqrt{\frac{2n!}{(n+L)!}} \zeta^{1/2+L} e^{-\kappa^2 \zeta^2 / 2} L_n^L(\kappa^2 \zeta^2)$$



- Eigenvalues

$$\mathcal{M}_{n,L,S}^2 = 4\kappa^2 \left(n + L + \frac{S}{2} \right)$$


$$4\kappa^2 \text{ for } \Delta n = 1$$

$$4\kappa^2 \text{ for } \Delta L = 1$$

$$2\kappa^2 \text{ for } \Delta S = 1$$




Orbital and radial states: $\langle \zeta \rangle$ increase with L and n



Parent and daughter Regge trajectories for the $I = 1$ ρ -meson family (red) and the $I = 0$ ω -meson family (black) for $\kappa = 0.54$ GeV

5 Higher-Spin Fermionic Modes

Hard-Wall Model



From Nick Evans

- Action for massive fermionic modes on AdS_{d+1} :

$$S[\bar{\Psi}, \Psi] = \int d^d x dz \sqrt{g} \bar{\Psi}(x, z) \left(i\Gamma^\ell D_\ell - \mu \right) \Psi(x, z)$$

- Equation of motion: $(i\Gamma^\ell D_\ell - \mu) \Psi(x, z) = 0$

$$\left[i \left(z\eta^{\ell m} \Gamma_\ell \partial_m + \frac{d}{2} \Gamma_z \right) + \mu R \right] \Psi(x^\ell) = 0$$

- Solution ($\mu R = \nu + 1/2$, $d = 4$)

$$\Psi(z) = C z^{5/2} [J_\nu(z\mathcal{M})u_+ + J_{\nu+1}(z\mathcal{M})u_-]$$

- Hadronic mass spectrum determined from IR boundary conditions $\psi_\pm(z = 1/\Lambda_{\text{QCD}}) = 0$


$$\mathcal{M}^+ = \beta_{\nu,k} \Lambda_{\text{QCD}}, \quad \mathcal{M}^- = \beta_{\nu+1,k} \Lambda_{\text{QCD}}$$

with scale independent mass ratio

- Obtain spin- J mode $\Phi_{\mu_1 \dots \mu_{J-1/2}}$, $J > \frac{1}{2}$, with all indices along 3+1 from Ψ by shifting dimensions

SU(6)	S	L	Baryon State
56	$\frac{1}{2}$	0	$N_{\frac{1}{2}}^{1+}$ (939)
	$\frac{3}{2}$	0	$\Delta_{\frac{3}{2}}^{\frac{3}{2}+}$ (1232)
70	$\frac{1}{2}$	1	$N_{\frac{1}{2}}^{1-}$ (1535) $N_{\frac{1}{2}}^{\frac{3}{2}-}$ (1520)
	$\frac{3}{2}$	1	$N_{\frac{1}{2}}^{\frac{1}{2}-}$ (1650) $N_{\frac{1}{2}}^{\frac{3}{2}-}$ (1700) $N_{\frac{1}{2}}^{\frac{5}{2}-}$ (1675)
	$\frac{1}{2}$	1	$\Delta_{\frac{1}{2}}^{\frac{1}{2}-}$ (1620) $\Delta_{\frac{1}{2}}^{\frac{3}{2}-}$ (1700)
56	$\frac{1}{2}$	2	$N_{\frac{1}{2}}^{\frac{3}{2}+}$ (1720) $N_{\frac{1}{2}}^{\frac{5}{2}+}$ (1680)
	$\frac{3}{2}$	2	$\Delta_{\frac{1}{2}}^{\frac{1}{2}+}$ (1910) $\Delta_{\frac{1}{2}}^{\frac{3}{2}+}$ (1920) $\Delta_{\frac{1}{2}}^{\frac{5}{2}+}$ (1905) $\Delta_{\frac{1}{2}}^{\frac{7}{2}+}$ (1950)
70	$\frac{1}{2}$	3	$N_{\frac{1}{2}}^{\frac{5}{2}-}$ $N_{\frac{1}{2}}^{\frac{7}{2}-}$
	$\frac{3}{2}$	3	$N_{\frac{1}{2}}^{\frac{3}{2}-}$ $N_{\frac{1}{2}}^{\frac{5}{2}-}$ $N_{\frac{1}{2}}^{\frac{7}{2}-}$ (2190) $N_{\frac{1}{2}}^{\frac{9}{2}-}$ (2250)
	$\frac{1}{2}$	3	$\Delta_{\frac{1}{2}}^{\frac{5}{2}-}$ (1930) $\Delta_{\frac{1}{2}}^{\frac{7}{2}-}$
56	$\frac{1}{2}$	4	$N_{\frac{1}{2}}^{\frac{7}{2}+}$ $N_{\frac{1}{2}}^{\frac{9}{2}+}$ (2220)
	$\frac{3}{2}$	4	$\Delta_{\frac{1}{2}}^{\frac{5}{2}+}$ $\Delta_{\frac{1}{2}}^{\frac{7}{2}+}$ $\Delta_{\frac{1}{2}}^{\frac{9}{2}+}$ $\Delta_{\frac{1}{2}}^{\frac{11}{2}+}$ (2420)
70	$\frac{1}{2}$	5	$N_{\frac{1}{2}}^{\frac{9}{2}-}$ $N_{\frac{1}{2}}^{\frac{11}{2}-}$ (2600)
	$\frac{3}{2}$	5	$N_{\frac{1}{2}}^{\frac{7}{2}-}$ $N_{\frac{1}{2}}^{\frac{9}{2}-}$ $N_{\frac{1}{2}}^{\frac{11}{2}-}$ $N_{\frac{1}{2}}^{\frac{13}{2}-}$

- Excitation spectrum for baryons in the hard-wall model: $\mathcal{M} \sim L + 2n$



Light baryon orbital spectrum for $\Lambda_{QCD} = 0.25$ GeV in the HW model. The **56** trajectory corresponds to L even $P = +$ states, and the **70** to L odd $P = -$ states: (a) $I = 1/2$ and (b) $I = 3/2$

Soft-Wall Model

- Equivalent to Dirac equation in presence of a holographic linear confining potential

$$\left[i \left(z \eta^{\ell m} \Gamma_{\ell} \partial_m + \frac{d}{2} \Gamma_z \right) + \mu R + \kappa^2 z \right] \Psi(x^{\ell}) = 0.$$

- Solution ($\mu R = \nu + 1/2$, $d = 4$)

$$\begin{aligned} \Psi_+(z) &\sim z^{\frac{5}{2}+\nu} e^{-\kappa^2 z^2/2} L_n^{\nu}(\kappa^2 z^2) \\ \Psi_-(z) &\sim z^{\frac{7}{2}+\nu} e^{-\kappa^2 z^2/2} L_n^{\nu+1}(\kappa^2 z^2) \end{aligned}$$


- Eigenvalues

$$\mathcal{M}^2 = 4\kappa^2(n + \nu + 1)$$

- Obtain spin- J mode $\Phi_{\mu_1 \dots \mu_{J-1/2}}$, $J > \frac{1}{2}$, with all indices along 3+1 from Ψ by shifting dimensions

$4\kappa^2$ for $\Delta n = 1$
 $4\kappa^2$ for $\Delta L = 1$
 $2\kappa^2$ for $\Delta S = 1$

\mathcal{M}^2



L

Parent and daughter **56** Regge trajectories for the N and Δ baryon families for $\kappa = 0.5$ GeV

6 Other Applications of Gauge/Gravity Duality to QCD






- Chiral symmetry breaking [Erlich, Katz, Son and Stephanov, Da Rold and Pomarol ...]
- Hadronic spectrum [Boschi-Filho, Braga, Frederico, Forkel, Beyer, Vega, Schmidt ...]
- Electromagnetic, gravitational and transition form-factors of composite hadrons
[Abidin and Carlson, Grigoryan and Radyushkin, Kwee and Lebed, Brodsky and GdT ...]
- DIS and Pomeron Physics [Polchinski, Strassler, Brower, Tan, Ballon Bayona, Boschi-Filho, Braga ...]
- Quark and gluon matter at extreme conditions in heavy ion physics (RHIC, LHC)
[Policastro, Son, Starinets, Kovtun, Gubser, Kim, Sin, Zahed, Cáceres, Güijosa, Edelstein, ...]
- Condensed matter physics [Herzog, Kovtun, Son ...]

Future Applications of Light-Front Holography

- Introduction of massive quarks (heavy and heavy-light quark systems)
- Systematic improvement (QCD Coulomb forces, higher Fock states (HFS) ...)

Example: Space- and Time Like Pion Form-Factor (HFS)

PRELIMINARY



$$|\pi\rangle = \psi_{q\bar{q}/\pi} |q\bar{q}\rangle + \psi_{q\bar{q}q\bar{q}/\pi} |q\bar{q}q\bar{q}\rangle$$

$$\mathcal{M}_\rho^2 \rightarrow 4\kappa^2(n + 1/2)$$

$$\kappa = 0.54 \text{ GeV}$$

$$\Gamma_\rho = 130, \Gamma_{\rho'} = 400, \Gamma_{\rho''} = 300 \text{ MeV}$$

$$P_{q\bar{q}q\bar{q}} = 13 \%$$