



AdS/QCD and Hadronic Phenomena



**Institute for Nuclear Theory Seminar
University of Washington
March 28, 2008**

*Stan Brodsky
SLAC, Stanford University*

Applications of AdS/CFT to QCD




Changes in physical length scale mapped to evolution in the 5th dimension z

in collaboration with Guy de Teramond

INT
March 28, 2008

AdS/QCD
2

Stan Brodsky, SLAC



8-2007
8685A14

- Truncated AdS/CFT (Hard-Wall) model: cut-off at $z_0 = 1/\Lambda_{\text{QCD}}$ breaks conformal invariance and allows the introduction of the QCD scale (Hard-Wall Model) [Polchinski and Strassler \(2001\)](#).
- Smooth cutoff: introduction of a background dilaton field $\varphi(z)$ – usual linear Regge dependence can be obtained (Soft-Wall Model) [Karch, Katz, Son and Stephanov \(2006\)](#).

We will consider both holographic models

Goal:

- **Use AdS/CFT to provide an approximate, covariant, and analytic model of hadron structure with confinement at large distances, conformal behavior at short distances**
- **Analogous to the Schrodinger Theory for Atomic Physics**
- *AdS/QCD Holographic Model*
- *Hadronic Spectra and Wavefunctions*

String Theory



AdS/CFT

Mapping of Poincare' and
Conformal $SO(4,2)$ symmetries of 3
+1 space
to AdS_5 space

Goal: First Approximant to QCD

Counting rules for Hard
Exclusive Scattering
Regge Trajectories
QCD at the Amplitude Level

AdS/QCD

Conformal Invariance +
Confinement at large
distances

Semi-Classical QCD / Wave Equations

Holography

Boost Invariant 3+1 Light-Front Wave Equations



$J=0, 1, 1/2, 3/2$ plus L

Integrable!

Hadron Spectra, Wavefunctions, Dynamics

AdS/QCD

Spacelike pion form factor from AdS/CFT



Data Compilation from Baldini, Kloe and Volmer



Soft Wall: Harmonic Oscillator Confinement



Hard Wall: Truncated Space Confinement


One parameter - set by pion decay constant.

de Teramond, sjb
See also: Radyushkin

AdS/QCD


Dirac's Amazing Idea: The Front Form

Evolve in
ordinary time



$$\sigma = ct - z$$

Evolve in
light-front time!



$$\tau = t + z/c$$

Instant Form

INT
March 28, 2008

AdS/QCD

Front Form


Stan Brodsky, SLAC

Light-Front Wavefunctions: rigorous representation of composite systems in quantum field theory

$$x = \frac{k^+}{P^+} = \frac{k^0 + k^3}{P^0 + P^3}$$

$$P^+, \vec{P}_\perp$$

Fixed $\tau = t + z/c$




$$\Psi_n(x_i, \vec{k}_{\perp i}, \lambda_i)$$

$$\sum_i^n x_i = 1$$

$$\sum_i^n \vec{k}_{\perp i} = \vec{0}_\perp$$

Invariant under boosts! Independent of P^μ

A Unified Description of Hadron Structure



Angular Momentum on the Light-Front

$$J^z = \sum_{i=1}^n s_i^z + \sum_{j=1}^{n-1} l_j^z.$$

Conserved
LF Fock state by Fock State

$$l_j^z = -i \left(k_j^1 \frac{\partial}{\partial k_j^2} - k_j^2 \frac{\partial}{\partial k_j^1} \right)$$

n-1 orbital angular momenta

Nonzero Anomalous Moment \rightarrow Nonzero orbital angular momentum

INT
March 28, 2008

AdS/QCD
10

Stan Brodsky, SLAC

$LF(3+1)$

AdS_5

$$\psi(x, \vec{b}_\perp) \quad \longleftrightarrow \quad \phi(z)$$

$$\zeta = \sqrt{x(1-x)\vec{b}_\perp^2} \quad \longleftrightarrow \quad z$$
$$\psi(x, \vec{b}_\perp) \rightarrow \text{blue oval} \quad \text{blue oval} \rightarrow z$$
$$x \quad (1-x)$$
$$\psi(x, \zeta) = \sqrt{x(1-x)} \zeta^{-1/2} \phi(\zeta)$$

Holography: Unique mapping derived from equality of LF and
AdS formula for current matrix elements: **em and gravitational!**

Holography: Map AdS/CFT to 3+1 LF Theory


Relativistic LF radial equation

Frame Independent

$$\left[-\frac{d^2}{d\zeta^2} + V(\zeta) \right] \phi(\zeta) = M^2 \phi(\zeta)$$

$$\zeta^2 = x(1-x)b_\perp^2.$$

G. de Teramond, sjb




Effective conformal potential:

$$V(\zeta) = -\frac{1-4L^2}{4\zeta^2} + \kappa^4 \zeta^2$$

confining potential:

Prediction from AdS/CFT: Meson LFWF



de Teramond, sjb

**“Soft Wall”
model**

$\kappa = 0.375$ GeV

massless quarks


$$\psi_M(x, k_\perp) = \frac{4\pi}{\kappa \sqrt{x(1-x)}} e^{-\frac{k_\perp^2}{2\kappa^2 x(1-x)}} \quad \phi_M(x, Q_0) \propto \sqrt{x(1-x)}$$

Hadron Distribution Amplitudes

Lepage, sjb

$$\phi_H(x_i, Q)$$

$$\sum_i x_i = 1$$



$$k_\perp^2 < Q^2$$

- Fundamental gauge invariant non-perturbative input to hard exclusive processes, heavy hadron decays. Defined for Mesons, Baryons

- Evolution Equations from PQCD,
OPE, Conformal Invariance

Lepage, sjb

Frishman, Lepage, Sachrajda, sjb

Peskin Braun

Efremov, Radyushkin Chernyak et al

- Compute from valence light-front wavefunction in light-cone gauge

$$\phi_M(x, Q) = \int^Q d^2 \vec{k} \psi_{q\bar{q}}(x, \vec{k}_\perp)$$

Light-Front QCD

Heisenberg Matrix Formulation

$$L^{QCD} \rightarrow H_{LF}^{QCD}$$


Physical gauge: $A^+ = 0$

$$H_{LF}^{QCD} = \sum_i \left[\frac{m^2 + k_\perp^2}{x} \right]_i + H_{LF}^{int}$$

H_{LF}^{int} : Matrix in Fock Space

$$H_{LF}^{QCD} |\Psi_h\rangle = \mathcal{M}_h^2 |\Psi_h\rangle$$

Eigenvalues and Eigensolutions give Hadron Spectrum and Light-Front wavefunctions



DLCQ: Periodic BC in x^- . Discrete k^+ ; frame-independent truncation

Light-Front Wave Functions in QCD

- Hadronic bound state expanded in n-particle Fock eigenstates $|\psi_h\rangle = \sum_n \psi_{n/h} |n\rangle$: the LF Hamiltonian $H_{LF} = P^2 = P^+ P^- - \mathbf{P}_\perp^2$, $H_{LF}|P\rangle = \mathcal{M}^2|P\rangle$, at fixed LF time $\tau = t + z/c$ (Dirac '49; Pauli and Pinsky, sjb Phys. Rept. 1988).
- Fock components

$$\psi_{n/h}(x_i, \mathbf{k}_{\perp i}) = \langle n; x_i, \mathbf{k}_{\perp i}, |\psi_h(P^+, \mathbf{P}_\perp)\rangle,$$

frame independent and encode hadron properties in high momentum-transfer collisions.

- Momentum fraction $x_i = k_i^+/P^+$ and $\mathbf{k}_{\perp i}$ are the relative coordinates of parton i in Fock-state n

$$\sum_{i=1}^n x_i = 1 \quad \sum_{i=1}^n \mathbf{k}_{\perp i} = 0.$$

- Define transverse position coordinates $x_i \mathbf{r}_{\perp i} = x_i \mathbf{R}_\perp + \mathbf{b}_{\perp i}$

$$\sum_{i=1}^n \mathbf{b}_{\perp i} = 0, \quad \sum_{i=1}^n x_i \mathbf{r}_{\perp i} = \mathbf{R}_\perp.$$

Light-Front QCD

Heisenberg Matrix Formulation

$$H_{LF}^{QCD} |\Psi_h\rangle = \mathcal{M}_h^2 |\Psi_h\rangle$$

DLCQ

Discretized Light-Cone Quantization

n	Sector	1 q̄q	2 gg	3 q̄q g	4 q̄q q̄q	5 gg g	6 q̄q gg	7 q̄q q̄q g	8 q̄q q̄q q̄q	9 gg gg	10 q̄q gg g	11 q̄q q̄q gg	12 q̄q q̄q q̄q g	13 q̄q q̄q q̄q q̄q
1	q̄q				
2	gg		
3	q̄q g	
4	q̄q q̄q	
5	gg g
6	q̄q gg							
7	q̄q q̄q g
8	q̄q q̄q q̄q	
9	gg gg
10	q̄q gg g
11	q̄q q̄q gg
12	q̄q q̄q q̄q g
13	q̄q q̄q q̄q q̄q	


Eigenvalues and Eigensolutions give Hadron Spectrum and Light-Front wavefunctions

H.C. Pauli & sjb

DLCQ: Frame-independent, No fermion doubling; Minkowski Space

LIGHT-FRONT SCHRODINGER EQUATION


$$\left(M_\pi^2 - \sum_i \frac{\vec{k}_{\perp i}^2 + m_i^2}{x_i} \right) \begin{bmatrix} \psi_{q\bar{q}/\pi} \\ \psi_{q\bar{q}g/\pi} \\ \vdots \end{bmatrix} = \begin{bmatrix} \langle q\bar{q} | V | q\bar{q} \rangle & \langle q\bar{q} | V | q\bar{q}g \rangle & \cdots \\ \langle q\bar{q}g | V | q\bar{q} \rangle & \langle q\bar{q}g | V | q\bar{q}g \rangle & \cdots \\ \vdots & \vdots & \ddots \end{bmatrix} \begin{bmatrix} \psi_{q\bar{q}/\pi} \\ \psi_{q\bar{q}g/\pi} \\ \vdots \end{bmatrix}$$



$$A^+ = 0$$


G.P. Lepage, sjb

Hadronization at the Amplitude Level




Construct helicity amplitude using Light-Front Perturbation theory; coalesce quarks via LFWFs

Light-Front Wavefunctions




Invariant under boosts! Independent of P^μ

Calculation of Form Factors in Equal-Time Theory



Need vacuum fluctuations

Calculation of Form Factors in Light-Front Theory




zero for $q^+ = 0$ **zero !!**

$$\frac{F_2(q^2)}{2M} = \sum_a \int [dx][d^2\mathbf{k}_\perp] \sum_j e_j \frac{1}{2} \times \quad \text{Drell, sjb}$$

$$\left[-\frac{1}{q^L} \psi_a^{\uparrow*}(x_i, \mathbf{k}'_{\perp i}, \lambda_i) \psi_a^{\downarrow}(x_i, \mathbf{k}_{\perp i}, \lambda_i) + \frac{1}{q^R} \psi_a^{\downarrow*}(x_i, \mathbf{k}'_{\perp i}, \lambda_i) \psi_a^{\uparrow}(x_i, \mathbf{k}_{\perp i}, \lambda_i) \right]$$


$$\mathbf{k}'_{\perp i} = \mathbf{k}_{\perp i} - x_i \mathbf{q}_\perp \quad \mathbf{k}'_{\perp j} = \mathbf{k}_{\perp j} + (1 - x_j) \mathbf{q}_\perp$$



Must have $\Delta \ell_z = \pm 1$ to have nonzero $F_2(q^2)$

Anomalous gravitomagnetic moment $B(0)$

Okun, Kobzarev, Teryaev: $B(0)$ Must vanish because of Equivalence Theorem




Hwang, Schmidt, sjb;
Holstein et al

$$B(0) = 0$$

Each Fock State

Hwang, sjb



- B decay amplitudes given by exact LF overlap formula
- Iterate interaction kernel whenever hard exchange occurs
- Factorization formulae at leading twist:
- Effective field theory approaches
- Key non-pert input: distribution amplitudes
- Fix renormalization scale: BLM
- Intrinsic charm Fock states contribute to annihilation amplitude, penguins

$$\mathcal{M} = \int \phi_B \times T_H \times \phi_D$$

Henley, Szczepaniak, sjb

Keum, Li, Sanda

Buchalla, Beneke, Neubert, Sachrajda

Bauer, Pirjol, Rothstein, Stewart

Lepage, sjb *Hoang*


Lepage, Mackenzie, sjb

Gardner, sjb

GPDs & Deeply Virtual Exclusive Processes

- New Insight into Nucleon Structure

Deeply Virtual Compton Scattering (DVCS)



x - quark momentum fraction

ξ - longitudinal momentum transfer


$\sqrt{-t}$ - Fourier conjugate to transverse impact parameter

$H(x, \xi, t), E(x, \xi, t), \dots$ “Generalized Parton Distributions”

Quark angular momentum (Ji sum rule)

$$J^q = \frac{1}{2} - J^G = \frac{1}{2} \int_{-1}^1 x dx [H^q(x, \xi, 0) + E^q(x, \xi, 0)]$$

X. Ji, Phys. Rev. Lett. 78, 610 (1997)



Light-cone wavefunction representation of deeply virtual Compton scattering \star

$$A_{J=0} \sim e_q^2 s^0 F(t)$$

*Local $J=0$
fixed pole
contribution*
Szczepaniak, Llanes-
Estrada, sjb

Link to DIS and Elastic Form Factors

DIS at $\xi=t=0$

$$H^q(x,0,0) = q(x), \quad -\bar{q}(-x)$$


$$\tilde{H}^q(x,0,0) = \Delta q(x), \quad \Delta \bar{q}(-x)$$

Form factors (sum rules)

$$\int_0^1 dx \sum_q [H^q(x, \xi, t)] = F_1(t) \text{ Dirac f.f.}$$

$$\int_0^1 dx \sum_q [E^q(x, \xi, t)] = F_2(t) \text{ Pauli f.f.}$$

$$\int_{-1}^1 dx \tilde{H}^q(x, \xi, t) = G_{A,q}(-t), \quad \int_{-1}^1 dx \tilde{E}^q(x, \xi, t) = G_{P,q}(-t)$$



Verified using
LFWFs
Diehl, Hwang, sjb

Quark angular momentum (Ji's sum rule)

$$J^q = \frac{1}{2} - J^G = \frac{1}{2} \int_{-1}^1 x dx [H^q(x, \xi, 0) + E^q(x, \xi, 0)]$$

X. Ji, Phys. Rev. Lett. 78, 610 (1997)

Conformal Theories are invariant under the Poincare and conformal transformations with

$$M^{\mu\nu}, P^\mu, D, K^\mu,$$

the generators of $SO(4,2)$

$SO(4,2)$ has a mathematical representation on AdS_5

Scale Transformations

- Isomorphism of $SO(4, 2)$ of conformal QCD with the group of isometries of AdS space

$$ds^2 = \frac{R^2}{z^2} (\eta_{\mu\nu} dx^\mu dx^\nu - dz^2), \quad \text{invariant measure}$$


$x^\mu \rightarrow \lambda x^\mu, z \rightarrow \lambda z$, maps scale transformations into the holographic coordinate z .

- AdS mode in z is the extension of the hadron wf into the fifth dimension.
- Different values of z correspond to different scales at which the hadron is examined.

$$x^2 \rightarrow \lambda^2 x^2, \quad z \rightarrow \lambda z.$$

$x^2 = x_\mu x^\mu$: invariant separation between quarks


- The AdS boundary at $z \rightarrow 0$ correspond to the $Q \rightarrow \infty$, UV zero separation limit.



INT
March 28, 2008

AdS/QCD
30


Stan Brodsky, SLAC



INT
March 28, 2008

AdS/QCD
31


Stan Brodsky, SLAC



INT
March 28, 2008

AdS/QCD
32

Stan Brodsky, SLAC



INT
March 28, 2008

AdS/QCD
33

Stan Brodsky, SLAC

AdS/CFT: Anti-de Sitter Space / Conformal Field Theory


Maldacena:

Map $AdS_5 \times S_5$ to conformal $N=4$ SUSY

- **QCD is not conformal;** however, it has manifestations of a scale-invariant theory:
Bjorken scaling, dimensional counting for hard exclusive processes
- **Conformal window:** $\alpha_s(Q^2) \simeq \text{const}$ at small Q^2
- **Use mathematical mapping of the conformal group $SO(4,2)$ to AdS₅ space**


Conformal QCD Window in Exclusive Processes


- Does α_s develop an IR fixed point? Dyson–Schwinger Equation Alkofer, Fischer, LLanes-Estrada, Deur ...
- Recent lattice simulations: evidence that α_s becomes constant and is not small in the infrared Furui and Nakajima, hep-lat/0612009 (Green dashed curve: DSE).



Deur, Korsch, et al: Effective Charge from Bjorken Sum Rule

$$\Gamma_{bj}^{p-n}(Q^2) \equiv \frac{g_A}{6} \left[1 - \frac{\alpha_s^{g_1}(Q^2)}{\pi} \right]$$






IR Fixed-Point for QCD?


- Dyson-Schwinger Analysis: QCD Coupling has IR Fixed Point
- Evidence from Lattice Gauge Theory
- Define coupling from observable: **indications of IR fixed point for QCD effective charges**
- Confined gluons and quarks have maximum wavelength: **Decoupling of QCD vacuum polarization at small Q^2** Serber-Uehling

$$\Pi(Q^2) \rightarrow \frac{\alpha}{15\pi} \frac{Q^2}{m^2} \quad Q^2 \ll 4m^2$$



- Justifies application of AdS/CFT in strong-coupling conformal window

Constituent Counting Rules



$$n_{tot} = n_A + n_B + n_C + n_D$$

Fixed t/s or $\cos \theta_{cm}$

$$\frac{d\sigma}{dt}(s, t) = \frac{F(\theta_{cm})}{s^{[n_{tot}-2]}} \quad s = E_{cm}^2$$


$$F_H(Q^2) \sim [\frac{1}{Q^2}]^{n_H-1}$$

Farrar & sjb; Matveev, Muradyan,
Tavkhelidze

Conformal symmetry and PQCD predict leading-twist scaling behavior of fixed-CM angle exclusive amplitudes

Characteristic scale of QCD: 300 MeV

Many new *J-PARC, GSI, J-Lab, Belle, Babar* tests



$$F_1(Q^2) \sim [1/Q^2]^{n-1}, \quad n = 3$$

From: M. Diehl *et al.* Eur. Phys. J. C **39**, 1 (2005).

- Phenomenological success of dimensional scaling laws for exclusive processes


$$d\sigma/dt \sim 1/s^{n-2}, \quad n = n_A + n_B + n_C + n_D,$$

implies QCD is a strongly coupled conformal theory at moderate but not asymptotic energies

Farrar and sjb (1973); Matveev *et al.* (1973).

- Derivation of counting rules for gauge theories with mass gap dual to string theories in warped space (hard behavior instead of soft behavior characteristic of strings) Polchinski and Strassler (2001).

Conformal behavior: $Q^2 F_\pi(Q^2) \rightarrow \text{const}$




Determination of the Charged Pion Form Factor at
 $Q^2=1.60$ and 2.45 (GeV/c^2).
 By Fpi2 Collaboration ([T. Horn et al.](#)). Jul 2006. 4pp.
 e-Print Archive: [nucl-ex/0607005](#)

Test of PQCD Scaling

Constituent counting rules

Farrar, sjb; Muradyan, Matveev, Tavkelidze



$s^7 d\sigma/dt(\gamma p \rightarrow \pi^+ n) \sim const$
fixed θ_{CM} scaling

PQCD and AdS/CFT:

$$s^{n_{tot}-2} \frac{d\sigma}{dt}(A + B \rightarrow C + D) = F_{A+B \rightarrow C+D}(\theta_{CM})$$

$$s^7 \frac{d\sigma}{dt}(\gamma p \rightarrow \pi^+ n) = F(\theta_{CM})$$

$$n_{tot} = 1 + 3 + 2 + 3 = 9$$



No sign of running coupling

Conformal invariance

INT
March 28, 2008

AdS/QCD
42

Stan Brodsky, SLAC




Conformal Invariance:

$$\frac{d\sigma}{dt}(\gamma p \rightarrow MB) = \frac{F(\theta_{cm})}{s^7}$$

Quark-Counting: $\frac{d\sigma}{dt}(pp \rightarrow pp) = \frac{F(\theta_{CM})}{s^{10}}$

$$n = 4 \times 3 - 2 = 10$$

P.V. LANDSHOFF and J.C. POLKINGHORNE



Angular distribution -- quark interchange

Best Fit

$$n = 9.7 \pm 0.5$$

Reflects underlying conformal scale-free interactions

INT
March 28, 2008

AdS/QCD
44

Stan Brodsky, SLAC