

Light-Front Holographic Approach to Strongly Coupled QCD


Guy F. de Téramond

Universidad de Costa Rica

Theoretical Physics Group

SLAC

March 9, 2011



1 Introduction

Gauge/Gravity Correspondence and Light-Front QCD

2 Light Front Dynamics

Light-Front Fock Representation

Semiclassical Approximation to QCD in the Light Front

3 Light-Front Holographic Mapping

Higher Spin Modes in AdS Space

Dual QCD Light-Front Wave Equation

AdS Bosonic Modes and Meson Spectrum

AdS Fermionic Modes and Baryon Spectrum

4 Light-Front Holographic Mapping of Current Matrix Elements


Electromagnetic Form Factors

Gravitational or Energy-Momentum Form Factor

1 Introduction

Gauge Gravity Correspondence and Light-Front QCD

- The AdS/CFT correspondence [Maldacena (1998)] between gravity on AdS space and conformal field theories in physical spacetime has led to a semiclassical approximation for strongly-coupled QCD, which provides physical insights into its non-perturbative dynamics
- Light-front (LF) quantization is the ideal framework to describe hadronic structure in terms of quarks and gluons: simple vacuum structure allows unambiguous definition of the partonic content of a hadron, exact formulae for form factors, physics of angular momentum of constituents ...
- Light-front holography provides a remarkable connection between the equations of motion in AdS and the bound-state LF Hamiltonian equation in QCD [GdT and S. J. Brodsky, PRL **102**, 081601 (2009)]
- Isomorphism of $SO(4, 2)$ group of conformal transformations with generators $P^\mu, M^{\mu\nu}, K^\mu, D$, with the group of isometries of AdS_5 , a space of maximal symmetry, negative curvature and a four-dim boundary: Minkowski space (Dim isometry group of AdS_{d+1} is $(d + 1)(d + 2)/2$)




2-2011
8685A14

- AdS₅ metric:

$$\underbrace{ds^2}_{L_{\text{AdS}}} = \frac{R^2}{z^2} \left(\underbrace{\eta_{\mu\nu} dx^\mu dx^\nu}_{L_{\text{Minkowski}}} - dz^2 \right)$$

- A distance L_{AdS} shrinks by a warp factor z/R as observed in Minkowski space ($dz = 0$):

$$L_{\text{Minkowski}} \sim \frac{z}{R} L_{\text{AdS}}$$




- Since the AdS metric is invariant under a dilatation of all coordinates $x^\mu \rightarrow \lambda x^\mu$, $z \rightarrow \lambda z$, the variable z acts like a scaling variable in Minkowski space
- Short distances $x_\mu x^\mu \rightarrow 0$ map to UV conformal AdS₅ boundary $z \rightarrow 0$
- Large confinement dimensions $x_\mu x^\mu \sim 1/\Lambda_{\text{QCD}}^2$ maps to large IR region of AdS₅, $z \sim 1/\Lambda_{\text{QCD}}$, thus there is a maximum separation of quarks and a maximum value of z
- Use the isometries of AdS to map the local interpolating operators at the UV boundary of AdS into the modes propagating inside AdS

2 Light Front Dynamics

- Different possibilities to parametrize space-time [Dirac (1949)]
- Parametrizations differ by the hypersurface on which the initial conditions are specified. Each evolve with different “times” and has its own Hamiltonian, but should give the same physical results
- *Instant form*: hypersurface defined by $t = 0$, the familiar one

- *Front form*: hypersurface is tangent to the light cone at $\tau = t + z/c = 0$



$$x^+ = x^0 + x^3 \quad \text{light-front time}$$


$$x^- = x^0 - x^3 \quad \text{longitudinal space variable}$$


$$k^+ = k^0 + k^3 \quad \text{longitudinal momentum} \quad (k^+ > 0)$$

$$k^- = k^0 - k^3 \quad \text{light-front energy}$$

$$k \cdot x = \frac{1}{2} (k^+ x^- + k^- x^+) - \mathbf{k}_\perp \cdot \mathbf{x}_\perp$$

On shell relation $k^2 = m^2$ leads to dispersion relation $k^- = \frac{\mathbf{k}_\perp^2 + m^2}{k^+}$





- Forms of Relativistic Dynamics: dynamical vs. kinematical generators [Dirac (1949)]

- Instant form

H, \mathbf{K} dynamical

\mathbf{L}, \mathbf{P} kinematical

- Point form

P^μ dynamical

$M^{\mu\nu}$ kinematical

- Front form

P^-, L^x, L^y dynamical

$P^+, \mathbf{P}_\perp, L^z, \mathbf{K}$ kinematical

- QCD Lagrangian

$$\mathcal{L}_{\text{QCD}} = -\frac{1}{4g^2} \text{Tr} (G^{\mu\nu} G_{\mu\nu}) + i\bar{\psi} D_\mu \gamma^\mu \psi + m\bar{\psi}\psi$$

- LF Momentum Generators $P = (P^+, P^-, \mathbf{P}_\perp)$ in terms of dynamical fields ψ, \mathbf{A}_\perp

$$P^- = \frac{1}{2} \int dx^- d^2 \mathbf{x}_\perp \bar{\psi} \gamma^+ \frac{(i\nabla_\perp)^2 + m^2}{i\partial^+} \psi + \text{interactions}$$

$$P^+ = \int dx^- d^2 \mathbf{x}_\perp \bar{\psi} \gamma^+ i\partial^+ \psi$$


$$\mathbf{P}_\perp = \frac{1}{2} \int dx^- d^2 \mathbf{x}_\perp \bar{\psi} \gamma^+ i\nabla_\perp \psi$$

- LF Hamiltonian P^- generates LF time translations

$$[\psi(x), P^-] = i \frac{\partial}{\partial x^+} \psi(x)$$

and the generators P^+ and \mathbf{P}_\perp are kinematical

Light-Front Fock Representation



- Dirac field ψ , expanded in terms of ladder operators on the initial surface

$$P^- = \sum_{\lambda} \int \frac{dq^+ d^2 \mathbf{q}_{\perp}}{(2\pi)^3} \left(\frac{\mathbf{q}_{\perp}^2 + m^2}{q^+} \right) b_{\lambda}^{\dagger}(q) b_{\lambda}(q) + \text{interactions}$$

- Construct LF Lorentz invariant Hamiltonian equation for the relativistic bound state

$$P_{\mu} P^{\mu} |\psi(P)\rangle = (P^- P^+ - \mathbf{P}_{\perp}^2) |\psi(P)\rangle = \mathcal{M}^2 |\psi(P)\rangle$$

- State $|\psi(P)\rangle$ is expanded in multi-particle Fock states $|n\rangle$ of the free LF Hamiltonian

$$|\psi\rangle = \sum_n \psi_n |n\rangle, \quad |n\rangle = \{ |uud\rangle, |uudg\rangle, |uud\bar{q}q\rangle, \dots \}$$

with $k_i^2 = m_i^2$, $k_i = (k_i^+, k_i^-, \mathbf{k}_{\perp i})$, for each constituent i in state n

- Fock components $\psi_n(x_i, \mathbf{k}_{\perp i}, \lambda_i^z)$ independent of P^+ and \mathbf{P}_{\perp} and depend only on relative partonic coordinates: momentum fraction $x_i = k_i^+ / P^+$, transverse momentum $\mathbf{k}_{\perp i}$ and spin λ_i^z

$$\sum_{i=1}^n x_i = 1, \quad \sum_{i=1}^n \mathbf{k}_{\perp i} = 0.$$

Semiclassical Approximation to QCD in the Light Front

[GdT and S. J. Brodsky, PRL **102**, 081601 (2009)]

- Compute \mathcal{M}^2 from hadronic matrix element $\langle \psi(P') | P_\mu P^\mu | \psi(P) \rangle = \mathcal{M}^2 \langle \psi(P') | \psi(P) \rangle$
- Find

$$\mathcal{M}^2 = \sum_n \int [dx_i] [d^2\mathbf{k}_{\perp i}] \sum_\ell \left(\frac{\mathbf{k}_{\perp \ell}^2 + m_\ell^2}{x_\ell} \right) |\psi_n(x_i, \mathbf{k}_{\perp i})|^2 + \text{interactions}$$

- LFWF ψ_n represents a bound state which is off the LF energy shell $\mathcal{M}^2 - \mathcal{M}_n^2$

$$\mathcal{M}_n^2 = \left(\sum_{a=1}^n k_a^\mu \right)^2 = \sum_a \frac{\mathbf{k}_{\perp a}^2 + m_a^2}{x_a}$$

with $k_a^2 = m_a^2$ for each constituent

- Invariant mass \mathcal{M}_n^2 key variable which controls the bound state: LFWF peaks at the minimum \mathcal{M}_n^2
- Semiclassical approximation to QCD:

$$\psi_n(k_1, k_2, \dots, k_n) \rightarrow \phi_n \left(\underbrace{(k_1 + k_2 + \dots + k_n)^2}_{\mathcal{M}_n^2} \right), \quad m_q \rightarrow 0$$

- In terms of $n - 1$ independent transverse impact coordinates $\mathbf{b}_{\perp j}, j = 1, 2, \dots, n - 1,$


$$\mathcal{M}^2 = \sum_n \prod_{j=1}^{n-1} \int dx_j d^2 \mathbf{b}_{\perp j} \psi_n^*(x_i, \mathbf{b}_{\perp i}) \sum_{\ell} \left(\frac{-\nabla_{\mathbf{b}_{\perp \ell}}^2 + m_{\ell}^2}{x_{\ell}} \right) \psi_n(x_i, \mathbf{b}_{\perp i}) + \text{interactions}$$

- Relevant variable conjugate to invariant mass in the limit of zero quark masses

$$\zeta = \sqrt{\frac{x}{1-x}} \left| \sum_{j=1}^{n-1} x_j \mathbf{b}_{\perp j} \right|$$

the x -weighted transverse impact coordinate of the spectator system (x active quark)

- For a two-parton system $\zeta^2 = x(1-x) \mathbf{b}_{\perp}^2$



- To first approximation LF dynamics depend only on the invariant variable ζ , and hadronic properties are encoded in the hadronic mode $\phi(\zeta)$ from

$$\psi(x, \zeta, \varphi) = e^{iM\varphi} X(x) \frac{\phi(\zeta)}{\sqrt{2\pi\zeta}}$$

factoring angular φ , longitudinal $X(x)$ and transverse mode $\phi(\zeta)$

- Ultra relativistic limit $m_q \rightarrow 0$ longitudinal modes $X(x)$ decouple ($L = L^z$)

$$\mathcal{M}^2 = \int d\zeta \phi^*(\zeta) \sqrt{\zeta} \left(-\frac{d^2}{d\zeta^2} - \frac{1}{\zeta} \frac{d}{d\zeta} + \frac{L^2}{\zeta^2} \right) \frac{\phi(\zeta)}{\sqrt{\zeta}} + \int d\zeta \phi^*(\zeta) U(\zeta) \phi(\zeta)$$

where the confining forces from the interaction terms are summed up in the effective potential $U(\zeta)$

- LF eigenvalue equation $P_\mu P^\mu |\phi\rangle = \mathcal{M}^2 |\phi\rangle$ is a LF wave equation for ϕ

$$\left(\underbrace{-\frac{d^2}{d\zeta^2} - \frac{1 - 4L^2}{4\zeta^2}}_{\text{kinetic energy of partons}} + \underbrace{U(\zeta)}_{\text{confinement}} \right) \phi(\zeta) = \mathcal{M}^2 \phi(\zeta)$$



- Effective light-front Schrödinger equation: relativistic, frame-independent and analytically tractable
- Eigenmodes $\phi(\zeta)$ determine the hadronic mass spectrum and represent the probability amplitude to find n -massless partons at transverse impact separation ζ within the hadron at equal light-front time
- Semiclassical approximation to light-front QCD does not account for particle creation and absorption but can be implemented in LF Hamiltonian EOM by applying the L-S formalism or evolution equations

3 Light-Front Holographic Mapping

Higher Spin Modes in AdS Space

- Description of higher spin modes in AdS space (Fronsdal, Fradkin and Vasiliev)
- Action for spin- J field in AdS_{d+1} in presence of dilaton background $\varphi(z)$ ($x^M = (x^\mu, z)$)

$$S = \frac{1}{2} \int d^d x dz \sqrt{g} e^{\varphi(z)} \left(g^{NN'} g^{M_1 M'_1} \dots g^{M_J M'_J} D_N \Phi_{M_1 \dots M_J} D_{N'} \Phi_{M'_1 \dots M'_J} - \mu^2 g^{M_1 M'_1} \dots g^{M_J M'_J} \Phi_{M_1 \dots M_J} \Phi_{M'_1 \dots M'_J} + \dots \right)$$

where D_M is the covariant derivative which includes parallel transport

$$[D_N, D_K] \Phi_{M_1 \dots M_J} = -R^L_{M_1 N K} \Phi_{L \dots M_J} - \dots - R^L_{M_J N K} \Phi_{M_1 \dots L}$$

- Physical hadron has plane-wave and polarization indices along $3+1$ physical coordinates

$$\Phi_P(x, z)_{\mu_1 \dots \mu_J} = e^{-iP \cdot x} \Phi(z)_{\mu_1 \dots \mu_J}, \quad \Phi_{z \mu_2 \dots \mu_J} = \dots = \Phi_{\mu_1 \mu_2 \dots z} = 0$$

with four-momentum P_μ and invariant hadronic mass $P_\mu P^\mu = M^2$

- Construct effective action in terms of spin- J modes Φ_J with only physical degrees of freedom
H. G. Dosch, S. J. Brodsky and GdT (in preparation)

- Introduce fields with tangent indices

$$\hat{\Phi}_{A_1 A_2 \dots A_J} = e_{A_1}^{M_1} e_{A_2}^{M_2} \dots e_{A_J}^{M_J} \Phi_{M_1 M_2 \dots M_J} = \left(\frac{z}{R}\right)^J \Phi_{A_1 A_2 \dots A_J}$$


- Find effective action

$$S = \frac{1}{2} \int d^d x dz \sqrt{g} e^{\varphi(z)} \left(g^{NN'} \eta^{\mu_1 \mu'_1} \dots \eta^{\mu_J \mu'_J} \partial_N \hat{\Phi}_{\mu_1 \dots \mu_J} \partial_{N'} \hat{\Phi}_{\mu'_1 \dots \mu'_J} - \mu^2 \eta^{\mu_1 \mu'_1} \dots \eta^{\mu_J \mu'_J} \hat{\Phi}_{\mu_1 \dots \mu_J} \hat{\Phi}_{\mu'_1 \dots \mu'_J} \right)$$

upon μ -rescaling

- Variation of the action gives AdS wave equation for spin- J mode $\Phi_J = \Phi_{\mu_1 \dots \mu_J}$

$$\left[-\frac{z^{d-1-2J}}{e^{\varphi(z)}} \partial_z \left(\frac{e^{\varphi(z)}}{z^{d-1-2J}} \partial_z \right) + \left(\frac{\mu R}{z} \right)^2 \right] \Phi_J(z) = \mathcal{M}^2 \Phi_J(z)$$



with $\hat{\Phi}_J(z) = (z/R)^J \Phi_J(z)$ and all indices along 3+1

Dual QCD Light-Front Wave Equation

$$\Phi_P(z) \Leftrightarrow |\psi(P)\rangle$$

- LF Holographic mapping found originally matching expressions of EM and gravitational form factors of hadrons in AdS and LF QCD [Brodsky and GdT (2006, 2008)]
- Upon substitution $z \rightarrow \zeta$ and $\phi_J(\zeta) \sim \zeta^{-3/2+J} e^{\varphi(z)/2} \Phi_J(\zeta)$ in AdS WE

$$\left[-\frac{z^{d-1-2J}}{e^{\varphi(z)}} \partial_z \left(\frac{e^{\varphi(z)}}{z^{d-1-2J}} \partial_z \right) + \left(\frac{\mu R}{z} \right)^2 \right] \Phi_J(z) = \mathcal{M}^2 \Phi_J(z)$$

find LFWE ($d = 4$)

$$\left(-\frac{d^2}{d\zeta^2} - \frac{1 - 4L^2}{4\zeta^2} + U(\zeta) \right) \phi_J(\zeta) = M^2 \phi_J(\zeta)$$

with

$$U(\zeta) = \frac{1}{2} \varphi''(z) + \frac{1}{4} \varphi'(z)^2 + \frac{2J - 3}{2z} \varphi'(z)$$

and $(\mu R)^2 = -(2 - J)^2 + L^2$

- AdS Breitenlohner-Freedman bound $(\mu R)^2 \geq -4$ equivalent to LF QM stability condition $L^2 \geq 0$
- Scaling dimension τ of AdS mode $\hat{\Phi}_J$ is $\tau = 2 + L$ in agreement with twist scaling dimension of a two parton bound state in QCD



Bosonic Modes and Meson Spectrum

- Positive dilaton background $\varphi = \kappa^2 z^2$: $U(z) = \kappa^4 \zeta^2 + 2\kappa^2(L + S - 1)$
- Normalized eigenfunctions $\langle \phi | \phi \rangle = \int d\zeta |\phi(z)|^2 = 1$

$$\phi_{nL}(\zeta) = \kappa^{1+L} \sqrt{\frac{2n!}{(n+L)!}} \zeta^{1/2+L} e^{-\kappa^2 \zeta^2 / 2} L_n^L(\kappa^2 \zeta^2)$$



- Eigenvalues

$$\mathcal{M}_{n,L,S}^2 = 4\kappa^2 (n + L + S/2)$$



LFWFs $\phi_{n,L}(\zeta)$ in physical spacetime for dilaton $\exp(\kappa^2 z^2)$: a) orbital modes and b) radial modes


$4\kappa^2$ for $\Delta n = 1$
 $4\kappa^2$ for $\Delta L = 1$
 $2\kappa^2$ for $\Delta S = 1$



Regge trajectories for the π ($\kappa = 0.6$ GeV) and the $I = 1$ ρ -meson and $I = 0$ ω -meson families ($\kappa = 0.54$ GeV)

Fermionic Modes and Baryon Spectrum

[Hard wall model: GdT and S. J. Brodsky, PRL **94**, 201601 (2005)]



From Nick Evans

- For baryons LFWE equivalent to system of coupled linear equations ($\nu = L + 1$)

$$-\frac{d}{d\zeta}\psi_- - \frac{\nu + \frac{1}{2}}{\zeta}\psi_- - \kappa^2\zeta\psi_- + 2i\kappa\psi_+ = \mathcal{M}\psi_+$$

$$\frac{d}{d\zeta}\psi_+ - \frac{\nu + \frac{1}{2}}{\zeta}\psi_+ - \kappa^2\zeta\psi_+ - 2i\kappa\psi_- = \mathcal{M}\psi_-$$

with eigenfunctions

$$\psi_+(\zeta) \sim \zeta^{\frac{1}{2}+\nu} e^{-\kappa^2\zeta^2/2} L_n^\nu(\kappa^2\zeta^2)$$

$$\psi_-(\zeta) \sim \zeta^{\frac{3}{2}+\nu} e^{-\kappa^2\zeta^2/2} L_n^{\nu+1}(\kappa^2\zeta^2)$$

and eigenvalues

$$\mathcal{M}^2 = 4\kappa^2(n + \nu)$$


- Large N_C : $\mathcal{M}^2 = 4\kappa^2(N_C + n + L - 2) \implies \mathcal{M} \sim \sqrt{N_C} \Lambda_{\text{QCD}}$

Same multiplicity of states for mesons and baryons!

$$4\kappa^2 \text{ for } \Delta n = 1$$

$$4\kappa^2 \text{ for } \Delta L = 1$$

$$2\kappa^2 \text{ for } \Delta S = 1$$



Regge trajectories for positive parity N and Δ baryon families ($\kappa = 0.5$ GeV)

4 Light-Front Holographic Mapping of Current Matrix Elements

[S. J. Brodsky and GdT, PRL **96**, 201601 (2006)], PRD **77**, 056007 (2008)]

- EM transition matrix element in QCD: local coupling to pointlike constituents

$$\langle \psi(P') | J^\mu | \psi(P) \rangle = (P + P')^\mu F(Q^2)$$

where $Q = P' - P$ and $J^\mu = e_q \bar{q} \gamma^\mu q$

- EM hadronic matrix element in AdS space from non-local coupling of external EM field propagating in AdS with extended mode $\Phi(x, z)$

$$\int d^4x dz \sqrt{g} e^{\varphi(z)} A^\ell(x, z) \Phi_{P'}^*(x, z) \overleftrightarrow{\partial}_\ell \Phi_P(x, z)$$

- Are the transition amplitudes related ?
- How to recover hard pointlike scattering at large Q out of soft collision of extended objects?

[Polchinski and Strassler (2002)]

- Mapping of J^+ elements at fixed light-front time: $\Phi_P(z) \Leftrightarrow |\psi(P)\rangle$

Electromagnetic Form-Factor

- Drell-Yan-West electromagnetic FF in impact space [Soper (1977)]

$$F(q^2) = \sum_n \prod_{j=1}^{n-1} \int dx_j d^2\mathbf{b}_{\perp j} \sum_q e_q \exp\left(i\mathbf{q}_{\perp} \cdot \sum_{k=1}^{n-1} x_k \mathbf{b}_{\perp k}\right) |\psi_n(x_j, \mathbf{b}_{\perp j})|^2$$

- Consider a two-quark π^+ Fock state $|u\bar{d}\rangle$ with $e_u = \frac{2}{3}$ and $e_{\bar{d}} = \frac{1}{3}$

$$F_{\pi^+}(q^2) = \int_0^1 dx \int d^2\mathbf{b}_{\perp} e^{i\mathbf{q}_{\perp} \cdot \mathbf{b}_{\perp}(1-x)} \left| \psi_{u\bar{d}/\pi}(x, \mathbf{b}_{\perp}) \right|^2$$

with normalization $F_{\pi^+}(q=0) = 1$

- Integrating over angle

$$F_{\pi^+}(q^2) = 2\pi \int_0^1 \frac{dx}{x(1-x)} \int \zeta d\zeta J_0\left(\zeta q \sqrt{\frac{1-x}{x}}\right) \left| \psi_{u\bar{d}/\pi}(x, \zeta) \right|^2$$

where $\zeta^2 = x(1-x)\mathbf{b}_{\perp}^2$

- Compare with electromagnetic FF in AdS space [Polchinski and Strassler (2002)]

$$F(Q^2) = R^3 \int \frac{dz}{z^3} V(Q, z) \Phi_{\pi^+}^2(z)$$

where $V(Q, z) = zQK_1(zQ)$

- Use the integral representation

$$V(Q, z) = \int_0^1 dx J_0 \left(\zeta Q \sqrt{\frac{1-x}{x}} \right)$$

- Find

$$F(Q^2) = R^3 \int_0^1 dx \int \frac{dz}{z^3} J_0 \left(zQ \sqrt{\frac{1-x}{x}} \right) \Phi_{\pi^+}^2(z)$$

- Compare with electromagnetic FF in LF QCD for arbitrary Q . Expressions can be matched only if LFWF is factorized



$$\psi(x, \zeta, \varphi) = e^{iM\varphi} X(x) \frac{\phi(\zeta)}{\sqrt{2\pi\zeta}}$$

- Find

$$X(x) = \sqrt{x(1-x)}, \quad \phi(\zeta) = \left(\frac{\zeta}{R} \right)^{-3/2} e^{\varphi(z)/2} \Phi(\zeta), \quad z \rightarrow \zeta$$

- “Free current” $V(Q, z) = zQK_1(zQ) \rightarrow$ infinite radius (mauve), no pole structure in time-like region
- “Dressed current” non-perturbative sum of an infinite number of terms \rightarrow finite radius (blue)
- Form factor in soft-wall model expressed as $N - 1$ product of poles along vector radial trajectory
[Brodsky and GdT (2008)] $(\mathcal{M}_\rho^2 \rightarrow 4\kappa^2(n + 1/2))$

$$F(Q^2) = \left[\left(1 + \frac{Q^2}{\mathcal{M}_\rho^2}\right) \left(1 + \frac{Q^2}{\mathcal{M}_{\rho'}^2}\right) \cdots \left(1 + \frac{Q^2}{\mathcal{M}_{\rho^{N-2}}^2}\right) \right]^{-1}$$





- Higher Fock components in pion LFWF

$$|\pi\rangle = \psi_{q\bar{q}/\pi}|q\bar{q}\rangle_{\tau=2} + \psi_{q\bar{q}q\bar{q}/\pi}|q\bar{q}q\bar{q}\rangle_{\tau=4} + \dots$$

- Expansion of LFWF up to twist 4 (monopole + tripole)

$$\kappa = 0.54 \text{ GeV}, \Gamma_\rho = 130, \Gamma_{\rho'} = 400, \Gamma_{\rho''} = 300 \text{ MeV}, P_{q\bar{q}q\bar{q}} = 13\%$$



Gravitational or Energy-Momentum Form-Factor

[S. J. Brodsky and GdT, PRD **78**, 025032 (2008)]

- Gravitational form factor of composite hadrons in QCD: local coupling to pointlike constituents

$$\langle P' | \Theta_{\mu}^{\nu} | P \rangle = (P^{\nu} P'_{\mu} + P_{\mu} P'^{\nu}) A(Q^2)$$




where $Q = P' - P$ and

$$\Theta_{\mu\nu} = \frac{1}{2} \bar{q} i (\gamma_{\mu} D_{\nu} + \gamma_{\nu} D_{\mu}) q - g_{\mu\nu} \bar{q} (i \mathcal{D} - m) q - G_{\mu\lambda}^a G_{\nu}^{a\lambda} + \frac{1}{4} g_{\mu\nu} G_{\lambda\sigma}^a G^{a\lambda\sigma}$$

- Hadronic matrix element of energy-momentum tensor from perturbing the static AdS metric: non-local coupling of external graviton field propagating in AdS with extended mode $\Phi(x, z)$

$$\int d^4x dz \sqrt{g} h_{\ell m} \left(\partial^{\ell} \Phi_{P'}^* \partial^m \Phi_P + \partial^m \Phi_{P'}^* \partial^{\ell} \Phi_P \right)$$

- Are the transition amplitudes related ?
- Mapping of Θ^{++} elements at fixed LF time: Identical mapping $\Phi_P(z) \Leftrightarrow |\psi(P)\rangle$ as EM FF



“ Working with a front is a process that is unfamiliar to physicists. But still I feel that the mathematical simplification that it introduces is all-important. I consider the method to be promising and have recently been making an extensive study of it. It offers new opportunities, while the familiar instant form seems to be played out ”

P.A.M. Dirac (1977)