INSUFLON

WHAT IS INSUFLON?

An indwelling subcutaneous (soft plastic) catheter for frequent subcutaneous injection without actual needle sticking the child.



WHERE TO INSERT INSUFLON? At the regular subcutaneous injection sites.



- a. Insuflon is due to be changed Q5days & prn (document insuflon site change on Vascular Access Device Hx)
- b. Always find a site with most subcutaneous fat



HOW TO USE INSUFLON?

	Step-by-Step Procedure	Nursing Assessment & Documentation
1.	Assess insuflon site before injection	- Document site assessment <u>before injection</u> on flowsheet.
2.	a. Check medication to be given (5 rights)	Potential complications: bleeding at site, bruising, redness, infection at site, swelling, other discoloration, and hardening tissue texture at site * Pt still feels the burning sensation of the Lovenox though! Lovenox: - low molecular weight heparin (LMWH)
	b. Draw med up in 1mL syringe	- a.k.a. enoxaparin - for anticoagulation
	c. Use <u>only</u> 27-31 gauge & 1/8 th – 3/8 th inch long needle or TB syringe	Monitoring blood test: Anti Xa level To be drawn @:
3.	a. Wash handb. Alcohol swab the port of insulfon for 15sec	
4.	 a. Align needle with port of insuflon b. Inject med through insuflon slowly with no resistance c. Finish injection, hold insuflon and pull needle off * NO FLUSHING needed!!!!! 	 Needle has to go in straight for proper administration of med If there is resistance, withdraw needle and realign
5.	DO NOT RUB the site	May cause bruising and irritation
6.	Insuflon and site change	Every 5 days Document site assessment on FLOWSHEET See online LPCH Procedure: "Insuflon Indwelling Subcutaneous Catheter" for insertion instruction