

Appendix D: Perineal Rash Algorithm

Use only warm water and cloths on bottoms.
No diaper wipes please

**Reddened bottoms
With intact skin**

Very mild areas of redness:
May use moisturizers
(i.e. Aquaphor or Eucerin)
PRN

Moderate to severe
areas of redness:
Sensicare Cream

**Reddened bottom with open areas
(Denuded, weepy raw skin)**

If mild skin breakdown, do the following PRN:

- Apply Stomahesive Powder over affected area.
- Then, apply Sensicare Cream
- 3. Apply more Stomahesive Powder

If moderate skin breakdown with increased
open areas of skin, do the following PRN:

- Fill the open areas of skin with Stomahesive
Powder then apply 3M No-Sting Barrier;
let dry. (May repeat twice, this is called "crusting").
- 2. Apply Sensicare Cream

**Reddened bottom with maculopapular rash
With or without open areas
(denuded, weepy raw skin)**

Apply antifungal cream or powder (i.e.
Nystatin or Lotrimin; requires
MD/AHP order)

Then the following PRN:
Apply Stomahesive Powder over
affected area.
Then, apply Sensicare Cream
Apply more Stomahesive Powder

If no improvement or skin breakdown worsens,
Consult the WOCN Team, pager #18611