



## Use of Mental Stress Management/Relaxation Techniques

In the past week (even if it was NOT a typical week), how many **times** did you do mental stress management or relaxation techniques?

none \_\_\_\_\_ times

Describe the mental stress management technique(s) you used: \_\_\_\_\_

### Scoring

This is a single item. If the technique described is not a cognitive symptom management technique, code as "0". Cognitive techniques include imagery, prayer, meditation, or progressive muscle relaxation. Reading, watching TV, listening to music, etc, are not considered cognitive techniques. To score the item, categorize the number of times into an ordinal scale with these categories:

- 1 = None
- 2 = 1-7 times/week
- 3 = 8 or more times/week

This item can also be left as a continuous measure, using the actual times coded.

### Characteristics

Tested on 1,129 subjects with chronic disease. N=51 for test-retest.

No. of items	Observed Range	Mean	Standard Deviation	Internal Consistency Reliability	Test-Retest Reliability
1	1-3	1.28	.53	—	.66

### Source of Psychometric Data

Stanford Chronic Disease Self-Management Study. Psychometrics reported in: Lorig K, Stewart A, Ritter P, González V, Laurent D, & Lynch J, *Outcome Measures for Health Education and other Health Care Interventions*. Thousand Oaks CA: Sage Publications, 1996, p.25.

## Comments

Another way that you can use this scale is to separate out the use of cognitive and non-cognitive techniques that one uses. We have never used the scale in this way but it seems like a reasonable possible use.

## References

Lorig K, Stewart A, Ritter P, González V, Laurent D, & Lynch J, *Outcome Measures for Health Education and other Health Care Interventions*. Thousand Oaks CA: Sage Publications, 1996, p.25,38.

*This scale is free to use without permission*

### **Stanford Patient Education Research Center**

1000 Welch Road, Suite 204  
Palo Alto CA 94304  
(650) 723-7935  
(650) 725-9422 Fax  
self-management@stanford.edu  
<http://patienteducation.stanford.edu>

Funded by the National Institute of Nursing Research (NINR)