

ISOLATION QUICK REFERENCE GUIDE

Contact	Droplet	Airborne – must be immune	Airborne – N95 needed	Immunosuppressed-Protective Isolation
<ul style="list-style-type: none"> • MRSA (<i>mask if in respiratory tract</i>) • VRE • Adenovirus • Diarrhea • C. dif • Rotavirus • E coli 0157 • Enterovirus • Salmonella • Shigella, • Hepatitis A • Herpes zoster (<i>shingles, localized</i>) • Herpes simplex • Parainfluenza (<i>mask if coughing</i>) • RSV (<i>mask if productive cough</i>) • Lice • Scabies 	<ul style="list-style-type: none"> • Pertussis • Influenza A or B • MRSA if in respiratory • Neisseria meningitidis (<i>suspected or confirmed</i>) • Coxsackie • Bacterial meningitis (for 24h after effective antibiotic therapy) • RSV(<i>Droplet & Contact</i>) 	<ul style="list-style-type: none"> • Chicken pox • Disseminated herpes zoster (<i>shingles</i>) • Measles 	<ul style="list-style-type: none"> • TB • SARS • Avian influenza 	<ul style="list-style-type: none"> • Bone marrow transplant or stem cell transplant • Solid organ transplant

Note: Standard Precautions are required for CMV, HIV, hepatitis B and C, Aspergillosis